



## Editorial Team

### Editor in Chief

» [Leka Lutpiatina](#), (Scopus ID: 57204557330) Department of Medical Laboratory Technology  
Poltekkes Kemenkes Banjarmasin, Indonesia

---

### Managing Editor

» [Ratih Dewi Dwiyanti](#), (Scopus ID : 57204558206) Department of Medical Laboratory Technology  
Poltekkes Kemenkes Banjarmasin, Indonesia

---

» [Meilana Dharma Putra](#), (Scopus ID : 50262867100) Department of Lambung Mangkurat  
Banjarmasin University, Indonesia

---

» [Çiğdem Ünal Kantekin](#), Department of Anesthesiology, Faculty of Medicine, University of Bozok,  
Turkey

---

### Editorial Boards Regional Europe

» [Luisa Barzon](#), (Scopus ID: 6603688122) Department of Molecular Medicine, University of  
Padova, Italy

---

» [Ozhan Ozdemir](#), Department of Obstetrics and Gynecology, Ankara Numune Training and  
Research Hospital, Ankara, Turkey

---

## Editorial Boards Regional Africa

» [Bashir Lawal](#), (Scopus ID : 57191326259) Department of Biochemistry, Tropical Disease Research Unit, Federal University of Technology, P.M.B. 65, Minna, Nigeria, Nigeria

---

» [Festus Abiose Olajubu](#), (Scopus ID: 55542617600) Department of Microbiology, Adekunle Ajasin University, Akungba-Akoko, Nigeria

---

## Editorial Boards Regional Asia

» [Dinna Rakhmina](#), (Scopus ID: 57208653176) Department of Medical Laboratory Technology Poltekkes Kemenkes Banjarmasin, Indonesia

---

» [I Nyoman Jirna](#), Medical Laboratory Technology Poltekkes Kemenkes Denpasar, Indonesia

---

» [Kurnia Dwi Artanti](#), Department of Epidemiology, Faculty of Public Health, Universitas Airlangga, Indonesia

---

» [Indah Purwaningsih](#), Medical Laboratory Technology Poltekkes Kemenkes Pontianak, Indonesia

---

» [Mahdalena Mahdalena](#), (Scopus ID: 57207763549) Quality Assurance Unit, Poltekkes Kemenkes Banjarmasin, Indonesia, Indonesia

---

» [Maria Tuntun](#), Medical Laboratory Technology Poltekkes Kemenkes Tanjung Karang Bandar Lampung, Indonesia

---

» [Neni Oktiyani](#), (Researcher ID : A-6676-2019) Department of Medical Laboratory Technology Poltekkes Kemenkes Banjarmasin, Indonesia

---

» [Ricko Dharmadi Utama](#), (Scopus ID: 57224725107) Department of Medical Laboratory Technology, Poltekkes Kemenkes Banjarmasin, Indonesia, Indonesia

---

# Vol 5, No 2 (2019)

**December**

DOI: <http://dx.doi.org/10.31964/mltj.v0i0>

## Table of Contents

---

### Articles

[Acute Toxicity Test of Black Pomegranate Peel Extract \(Granati Fructus Cortex\) Against Larvae of Shrimp \(\*Artemia salina\* Leach\)](#)

[PDF](#)

DOI : [10.31964/mltj.v0i0.242](https://doi.org/10.31964/mltj.v0i0.242) | Abstract views : 456 times

62-69

Rofiatu Sholihah Achmad, Lingga Aditya Aditya, Nosa Ika Cahyariza

[Anti-hyperglycemic Effect of Torbangun \(\*Coleus amboinicus\* Lour\) Leaves Extract Through Liver and Muscle Glycogen Deposits in Streptozotocin-induced Hyperglycemic Sprague-Dawley Rats Model](#)

[PDF](#)

DOI : [10.31964/mltj.v0i0.227](https://doi.org/10.31964/mltj.v0i0.227) | Abstract views : 626 times

70-80

Meilla Dwi Andrestian, Rizal Damanik, Faisal Anwar, Nancy Dewi Yuliana

[Young and Old Sugar Apple \(\*Annona squamosa\* Linn\) Leaf Extracts As an \*Aedes aegypti\* Larva Insecticide](#)

[PDF](#)

DOI : [10.31964/mltj.v0i0.231](https://doi.org/10.31964/mltj.v0i0.231) | Abstract views : 377 times

81-87

Abdul Khair, Noraida Noraida

[Ethanol Extract of Black Cumin Seed \(Nigella Sativa\) Reduces Expression Kidney iNOS in Mice Model Preeclampsia](#)

[PDF](#)

DOI : [10.31964/mltj.v0i0.247](#) | Abstract views : 318 times

88-95

Rubiati Hipni

[Quality of Genom in Type II Diabetes Mellitus Patients in Viewed of Temperature, Storage Duration, Number of Leukocyte](#)

[PDF](#)

DOI : [10.31964/mltj.v0i0.229](#) | Abstract views : 337 times

96-101

Puspitasari Puspitasari, Evi Rinata, Rohman Dijaya, Siska Aprilia, Dina Trikumalasari, Livia Nur Azzah, Qilmia Fanani, Miftahul Mushlih, Andika Aliviameita, Dian Delta

[Detection of Salmonella typhi Using Multiplex and Monoplex PCR in Tifoid Fever Patients](#)

[PDF](#)

DOI : [10.31964/mltj.v0i0.230](#) | Abstract views : 412 times

108-107

Andika Aliviameita, Meryn Linanda Agustin, Anggie Wedha Puspita, Miftahul Mushlih, Puspitasari Puspitasari, Yanik Purwanti, Arief Wisaksono

[Analysis between Adrenal MRI Mass Size and 1 mg Dexamethasone Suppression Test in Patients with Adrenal Incidentaloma](#)

[PDF](#)

DOI : [10.31964/mltj.v0i0.223](#) | Abstract views : 377 times

108-116

Pinar Karakaya, Bulent Yaprak, Bahar Ozdemir, Hakan Kocoglu

[Analysis of Obesity And Axillary Lymph Nodes on Mammography](#)

[PDF](#)

DOI : [10.31964/mltj.v0i0.235](#) | Abstract views : 297 times

117-123

Aysegul Akdogan Gemici, Ersoy Bayram, Nigar Gultekin, Ercan Inci

Detection of Mycobacterium tuberculosis by Microscopic Technique of Sputum AFB and ICT (Cocktail Antigen) TB

[PDF](#)

DOI : [10.31964/mltj.v0i0.246](https://doi.org/10.31964/mltj.v0i0.246) | Abstract views : 453 times

124-  
130

Retno Martini Widhyasih, Annisa Husnun Hanifah, Chairlan Chairlan, Dewi Inderiati

Analysis between Platelet Count and Blood Groups in Apheresis Platelet Donors with Demographic Features

[PDF](#)

DOI : [10.31964/mltj.v0i0.250](https://doi.org/10.31964/mltj.v0i0.250) | Abstract views : 455 times

131-  
137

Canan Eren, Serpil Çeçen



## Young and Old Sugar Apple (*Annona squamosa* Linn) Leaf Extracts As an *Aedes aegypti* Larva Insecticide

\*Abdul Khair, Noraida

Environmental Health Poltekkes Kemenkes Banjarmasin  
Mistar Cokrokusumo Street 1a Banjarbaru Indonesia.

\*Email: [ulunkhair@gmail.com](mailto:ulunkhair@gmail.com)

DOI: 10.31964/mltj.v%vi%i.231

**Abstract:** Vector-borne diseases are still a health problem in some tropical countries. One vector-borne disease is Dengue Hemorrhagic Fever (DHF). DHF never decreases and even tends to continue to increase, and many cause deaths in children, 90% of them attack children under 15 years. DHF cases in 2018 amounted to 65,602 cases, with 467 deaths (CFR = 0.71%). The behavior of holding water in various places such as tubs and tendons indirectly creates a breeding place for *Aedes aegypti* mosquitoes. The use of chemicals as insecticides can cause the death of non-target animals, environmental pollution, and the occurrence of vector resistance to insecticides. Therefore it is necessary to do other methods including the use of vegetable insecticides. Sugar apple leaves can be used to kill *Aedes aegypti* larvae. Sugar apple leaves will obtain throughout the year. The purpose of this study is to know the ability of young and old sugar apple leaves to kill *Aedes aegypti* larvae. This type of research is a posttest only control group design. The study sample was part of the *Aedes aegypti* larvae on the final instar III. The experiment to kill *Aedes aegypti* larvae use two types of sugar apple leaf extract, namely young and old sugar apple leaves with nine treatments four replications. Each treatment consisted of 25 larvae. The research results show There was no significant difference in the number of dead larvae using either extracts from young sugar apple leaves or old sugar apple leaves. LC90 of sugar apple leaf extract was between 0.05632 to 0.08324% and the effective residual age at LC90 (0.06568%) with the death of *Aedes aegypti* larvae was 92% over 24 hours (1 day). **Keywords:** sugar apple; *Aedes aegypti*

### INTRODUCTION

Vector-borne diseases are still a health problem in some tropical countries. One vector-borne disease is Dengue Hemorrhagic Fever (DHF). DHF never decreases and even tends to continue to increase, and many cause deaths in children, 90% of them attack children under 15 years (Candra, 2009). The mosquito species that can be a DHF vector is *Aedes aegypti*. Since it first discovered in Surabaya and Jakarta in 1968, the number of dengue cases continues to spread to all regions in Indonesia. DHF cases in 2018 amounted to 65,602 cases, with 467 deaths (CFR = 0.71%)(Data and Information Center, 2015).

Residents usually store water in various places such as tubs and tendons. It estimated that each house has 5 to 6 reservoirs. The behavior of holding water is very dependent on the culture and water needs of the community. In areas where the piping system already exists, people still tend to store water as a backup if at any time the water does not flow. This habit indirectly creates a breeding place for *Aedes*

**Corresponding Author:** Abdul Khair

Environmental Health Poltekkes Kemenkes Banjarmasin

Mistar Cokrokusumo Street 1a Banjarbaru Indonesia Email: [ulunkhair@gmail.com](mailto:ulunkhair@gmail.com)

aegypti mosquitoes. The results of the study showed that of the 268 containers examined; there were 82 containers (30.6%) containing *Aedes aegypti* larvae. The value of the Container Index (CI) is 30.6, and the Breteau Index (BI) is 82 (Yudhastuti et al., 2005).

Until now, vector control is more focused on the use of chemicals because they are considered more effective, easy, and fast. However, the use of chemicals can cause the death of non-target animals, environmental pollution, and the occurrence of vector resistance to insecticides. Chemical DHF control includes using Temephos (Abate®) as larvicide and malathion as adulticides. Temephos usually used in water reservoirs because of their very low toxicity to humans. The use of 1 mg/liter Temephos can control the *Aedes aegypti* larvae in breeding sites for 3 to 3.5 months (Kurniawan et al., 2019). The use of Temephos 1% at a dose of 1 gr / 10 liters does not guarantee the permanent removal of larvae in breeding sites because people generally do not like the smell caused by larvicides. The use of Temephos must also be routine. Vector control by chemical means requires funds that are not small and harm the environment (Hasibuan, 2004). In three subdistricts (Tambaksari, Gubeng, and Sawahan), the mortality rates were under 80%, indicating possible resistance to Temephos (Mulyatno et al., 2012). Therefore it is necessary to do other methods including the use of vegetable insecticides.

In general, plant insecticides interpreted as insecticide whose essential ingredients come from plants. Because it made from natural materials, this type of pesticide is biodegradable, so it does not pollute the environment and is relatively safe for humans and livestock. Vegetable insecticides are hit and run if applied to kill pests at that time, and after the period executed, the residue will quickly decompose in nature. The power to kill plant-based insecticides is because there are toxic substances which are stomach poisons and contact poisons in soft-bodied animals. The use of vegetable insecticides is intended not to abandon and prohibit the use of synthetic pesticides, but only an alternative method with the aim that users do not depend solely on synthetic insecticides. Another goal is to minimize the use of synthetic insecticides so that the environmental damage caused can reduce. Besides that, the use of plant-based pesticides will improve agro-industrial development, especially rural industries, new business growth, and environmental sustainability.

In Indonesia, which is a tropical country, a variety of plants can grow well, and many of them are sugar apple (*Annona squamosa* Linn). In the form of extracts, sugar apple is used as antifungal and antioxidant activities (Kalidindi et al., 2015), inhibiting the growth of *Escherichia coli* bacteria and *Staphylococcus aureus* (Tansil et al., 2016), antihyperlipidemic in humans (Rofida et al., 2015), and to control leaf-eating pest on soybean (Indiati, 2014). Sugar apple also used as an insecticide. *Annona squamosa* seed extract can be used to control *Aedes albopictus* and *Culex quinquefasciatus* (Ravaomanarivo et al., 2014).

All parts of sugar apple plants can be used as insecticides (Mondal et al., 2018). Sugar apple which contains active ingredients as vegetable insecticides are raw fruit, leaves, and roots. The active ingredient works as a contact and stomach poison and is as an insecticide, repellent, and antifeedant. Sugar apple seed extract has been used to kill *Aedes aegypti* larvae (Wahyuni et al., 2015). Sugar apple seeds solution with liquid electric method had used as an insecticide against the mortality of *Aedes aegypti* (Novasari et al., 2017). But the use of sugar apple seeds cannot be obtained throughout the year. Sugar apple seeds are only available at certain times, from January to March.



In a plant, there are usually leaves. There are parts of leaves that are still young and pieces of leaves that are old. A comparison between young leaves and young leaves of plants is not only done to determine the effect of the active compound content but also on its activity. The results of the study of Felicia et al. (2016) note that old avocado leaves have higher antioxidant activity than young avocado leaves. The difference in the age of leaves of each plant is not always the same between young leaves and old leaves. However, the presence of higher levels of active compounds can provide equal or more significant activity between immature and mature leaves (Mulangsri, 2018).

In sugar apple plants, the leaves can also use as an insecticide. The advantage is that sugar apple leaves will obtain throughout the year. It can be used to kill *Aedes aegypti* larvae (Purwaningsih et al., 2015). There are young leaves and old leaves on the sugar apple tree. Furthermore, it is necessary to know the ability of young and old sugar apple leaves to kill *Aedes aegypti* larvae.

## **MATERIALS AND METHOD**

This type of research is a posttest only control group design. The study population was *Aedes aegypti* larvae on the final instar III. The study sample was part of the *Aedes aegypti* larvae on the final instar III. The experiment to kill *Aedes aegypti* larvae using two types of sugar apple leaf extract, namely young and old sugar apple leaves. Tests were carried out as many as nine treatments, both for attempting to kill larvae using young sugar apple leaves or using old sugar apple leaves. Each treatment carried out four replications. According to WHO standards for bioassay tests, each treatment consisted of 25 larvae (WHO, 2005).

Sugar apple leaf extract is made by maceration, which is soaked in ethanol solvent 70% for one night, and then the solution is taken. This treatment is carried out for three times to ensure that the leaves have extract properly. After that, the solvent evaporated with a rotary evaporator (RV10 IKA); the concentrated sugar apple leaf extract obtained.

Research procedures include carrying out preliminary experiments, determining LC50 and LC90, and determining the residual age. Initial tests were conducted to determine the types of sugar apple leaves and concentrations that could kill *Aedes aegypti* larvae. The determination of LC50 and LC90 was carried out to evaluate the effectiveness of larvacide toxicity through a bioassay test. Determination of residual age was carried out to determine the length of time the residue was still able to kill *Aedes aegypti* larvae. Data obtained by counting the number of larvae that died during the trial period. Descriptive data analysis is presented in the form of tables, while analytic data analysis is done using a t-test, ANOVA, and probit analysis.

## **RESULTS AND DISCUSSION**

### **Preliminary and Follow-up Test**

All treatment groups for both young and old sugar apple leaf extracts in the preliminary test showed the death of *Aedes aegypti* larvae. The higher the concentration of sugar apple leaf extracts given, the number of *Aedes aegypti* larvae that died. *Aedes aegypti* larvae did not die in the control and new groups. This proves that the death of *Aedes aegypti* larvae in the treatment group was caused by sugar apple leaf extract, not by the ethanol 70% or distilled water used in this study.



Table 1. Preliminary Test

Concentration	Death of Larvae			
	Young Sugar Apple Leaf Average	Young Sugar Apple Leaf (%)	Old Sugar Apple Leaf Average	Old Sugar Apple Leaf (%)
0.01563	2.3	9.2	2.6	10.4
0.03125	9	36	10.6	42.4
0.0625	21.7	86.8	22.4	89.6
0.125	24.3	97.2	24.7	98.8
0.25	25	100	25	100
0.5	25	100	25	100
1	25	100	25	100
Control (0.5 ml ethanol 70%+aquades)	0	0	0	0
Blank (aquades)	0	0	0	0

Table 2. Normality and Paired T Test (CI 95%)

	Normality Test	Paired T Test
	P	P
0.01563	0.707	0.519
0.03125	2.500	0.067
0.0625	2.121	0.101
0.125	0.707	0.519

The results of statistical analysis for all extract concentrations using Paired T-Test showed p-value > 0.05. That means that there is no significant difference between young and old sugar apple leaf extracts at all concentrations tested to kill *Aedes aegypti* larvae. This shows that the content of active compounds found in young and old sugar apple leaves is the same.

Based on the analysis of variance, the difference in the average number of *Aedes aegypti* larvae deaths was due to the effect of the concentration level of sugar apple leaf extract.

The results of statistical analysis using the Anova Test showed a p-value <0.05. That means that there are significant differences between the various concentrations of sugar apple leaf extract tested to kill *Aedes aegypti* larvae. Further analysis using the LSD method found that the level of sugar apple leaf extract of 0.0586% and 0.0676% showed the number of deaths of the same *Aedes aegypti* larvae. Thus it can be said that the concentration of 0.0586% has a sufficient level in influencing the mortality of *Aedes aegypti* larvae.

Table 3. Follow-up Test

Concentration	Death of Larvae	
	Average	%
0.0136 <sup>a</sup>	1	4
0.0226 <sup>b</sup>	5.25	21
0.0316 <sup>b</sup>	11.25	45
0.0406 <sup>c</sup>	15.75	63
0.0496 <sup>ca</sup>	18	72
0.0586 <sup>ca</sup>	20.75	83
0.0676 <sup>c</sup>	24	96
Control (0.5 ml ethanol 70%+aquades)	0	0
Blank (aquades)	0	0
Anova test	0.000	

**LC50 and LC90 Tests**

The lowest average of dead *Aedes aegypti* larvae occurred at sugar apple leaf extract concentration of 0.0136%, which is one larva (4%) and the highest average of dead *Aedes aegypti* larvae occurred in sugar apple leaf extract 0.0676%, which is 24 larvae (96%). So we can say that all concentrations of sugar apple leaf extract used in the LC50 and LC90 tests can cause death in *Aedes aegypti* larvae. The higher level of sugar apple leaf extract used will cause more *Aedes aegypti* larvae to die.

Table 4. LC50 and LC90 Tests

Death of Larvae (%)	Concentration	CI 95%
50	0.03418	0.03418 < LC < 0.03823
90	0.06568	0.05632 < LC < 0.08324

**Residual Age Test**

In the residual life test, old sugar apple leaf extract used at LC90 concentration of 0.06568% with the assumption that the level was considered very useful to kill *Aedes aegypti* larvae.

Table 5. Residual Age Test

Concentration of Extract (%)	Number of Larvae	Death of Larvae (%)							
		1	2	3	4	5	6	7	8
0.06568 (LC90)	25	96	86	74	61	43	26	2.2	0
Control	25	0	0	0	0	0	0	0	0
Blank	25	0	0	0	0	0	0	0	0

Based on the results of the residual life test obtained by larval mortality at two days (24 hours) remaining age of 92% with 24 hours contact time. This means that the old sugar apple leaf extract with a concentration of 0.06568% has a useful residue of more than 24 hours (1 day), although for the next residual age the larval mortality has decreased. This is because plant-based insecticides are "hit and run," if applied they will kill pests at that time and after their pests killed the residue will quickly decompose in nature.

Table 4 shows that LC50 and LC90 were 0.03418% and 0.06568%, respectively. The LC50 score in this study was slightly higher compared to the survey conducted by Purwaningsih et al. (2005) in the amount of 0.00902%.

In another study, it was found that LC50 from betel leaf extract was 910.38 or 0.09104% and LC50 from *Annona squamosa* seed extract was 20.62 ppm or 0.00206% (Wahyuni et al., 2015). This shows that the toxicity of sugar apple seed extract is still better when compared to sugar apple leaf extract in this study (0.03148%). However, the toxicity of sugar apple leaf extract is better than betel leaf extract.

Sugar apple leaves contain annonain which are effective active ingredients as vegetable insecticides, work as stomach poisons and contact poisons against insects. The toxic effects of a sugar apple seed or leaf extract contact can see from the clinical symptoms that arise in the larvae, namely the movement becomes slow, the body shrinks and eventually dies. These symptoms indicate that the larva is running out of energy (ATP) (Wardhana et al., 2005). Known annonain is known to act as a dissolved poison or contact poison. It can inhibit electron transfer by blocking the bond between NADH and ubiquinone in the electron transfer chain in the cell respiration process, causing the metabolic energy formation process to be inhibited (Aisyafahmi et al., 2017).

## CONCLUSION

There was no significant difference in the number of dead larvae using either extracts from young sugar apple leaves or old sugar apple leaves. LC90 of sugar apple leaf extract was between 0.05632 to 0.08324% and the sufficient residual age at LC90 (0.06568%) with the death of *Aedes aegypti* larvae was 92% over 24 hours (1 day).

## REFERENCE

- Aisyafahmi, D. and Wahyuni, D. (2017). Toxicity of Granule from Sugar Apple (*Annona squamosa*. L) Fruit Extract on The Mortality *Aedes aegypti* Larvae. *Bioedukasi*, XV(2), 26-30.
- Candra, A. (2009). Dengue Hemorrhagic Fever: Epidemiology, Pathogenesis, and Its Transmission Risk Factors. *Aspirator*, 2(2), 110-119.
- Data and Information Center (2019). *Indonesia Health Profile 2018*. Ministry of Health of the Republic of Indonesia.
- Felicia, N., Widarta, I.W.R. & Yusarini N.L.A. (2016). Effect of Leaf Leaves and Processing Methods on Antioxidant Activity and Sensory Characteristics of Herbal Tea Avocado Leaf Powder (*Persea americana* Mill.). *OJS Universitas Udayana*.
- Hasibuan, R. (2004). Field Evaluation of the Impact of Isoproc carb Insecticide Application in Predator Insect and Pest of *Aulacaspis tegalensis* Zhnt. (Homoptera: Diaspididae) in Sugarcane Plantations. *Tropical Plants Pests and Diseases Journal*, 4(2), 69-74.

- Indiati, S.W. (2014). The Use of Sugar Apple and Neem Extract to Control Leaf Eating Pest on Soybean. *Journal of Experimental Biology and Agricultural Sciences* 2(2), 208-2014.
- Kalidindi, N., Thimmaiah, N.V., Jagadeesh, N.V., Nandeeep, R., Swetha, S., & Kalidindi, B. (2015). Antifungal and Antioxidant Activities of Organic and Aqueous Extract of *Annona squamosa* Linn. Leaves. *Journal of Food and Drug Analysis*, 23(4), 795-802.
- Kurniawan, A., Nurjana, M.A., Srikandi, Y. (2019). The Use of Temephos in Household and its Effect Density of The *Aedes* sp Larvae in The Balaroa Lower District , Palu City. *Disease Vector Journal*, 13(1), 67-76.
- Mondal, P., Biswas, S., Pal, K., and Ray, D.P. (2018). *Annona squamosa* as a Potential Botanical Insecticide for Agricultural Domains: A Review. *International Journal of Bioresource Science*, 5(1), 81-89.
- Mulangsi, D.A.K. (2018). Antibacterial Activities of Ethanol Extract of Young Leaves and Green Betel (*Piper betle* L.) Leaves on *Staphylococcus aureus* Bacteria, *Jurnal Ilmiah Cendekia Eksakta*, 1-4
- Mulyatno, K.C., Yamanaka, A., Ngadino, & Konoshi, E. (2012). Resistance of *Aedes aegypti* (L.) Larvae to Temephos in Surabaya, Indonesia. *Southeast Asian Journal of Tropical Medicine and Public Health*, 43(1), 29-33.
- Novasari, A.M. & Sasongkowati, R. (2017). The Effect of Sugar Apple (*Annona squamosa* L.) Seeds Solution as an Insecticide against Mortality of *Aedes aegypti* Mosquito with Liquid Electric Method. *Environmental Health Journal* 9(2), 200-208.
- Purwaningsih, N.V., Kardiwinata, M.P., Utami, N.W.A. (2015). The Killing Power of Sugar Apple (*A. squomosa* L.) Leaf Extracts against Eggs and *A. aegypti* Larvae. *Cakra Kimia (Indonesia E-Journal of Applied Chemistry)*, 3(2), 96-103.
- Ravaomanarivo, L.H.R., Razafindraleva, H.A., Raharimalala, F.N., Rasoahantaveloniaina, B., Ravelonandro, P.H., & Mavingui, P. (2014). Efficacy of seed extracts of *Annona squamosa* and *Annona muricata* (Annonaceae) for the control of *Aedes albopictus* and *Culex quinquefasciatus* (Culicidae). *Asian Pacific Journal of Tropical Biomedicine*, 4(10), 798-806.
- Rofida, S., Firdiansyah, A., & Fitriyastuti, E. (2015). Antihyperlipidemia Activity of *Annona squamosa* L. Leaves Ethanol Extract *Journal of Pharmaceutical Science and Pharmacy Practice*, 2(1), 1-3.
- Tansil, A.Y.M., Nangoy, E., Posangi, J., Bara, R.A. (2016). Inhibition Test of Ethanol Extract of Sugar Apple Leaf (*Annona squamosa*) on the Growth of *Escherichia coli* and *Staphylococcus aureus* Bacteria. *e-Biomedik Journal (eBm)*, 4(2).
- Wahyuni, D., Loren, I. (2015). Difference between Toxicity of Betel Leaf Extract (*Piper betle* L.) and Sugar Apple (*Annona squamosa* L.) Seed Extract against *Aedes aegypti* Mosquito Larvae. *Jurnal Saintifika*, 17(1), 38-48.
- Wardhana A. H., A. Husein dan J. Manurung. 2005. The Effectiveness of Sugar Apple Seed Extract with Water, Methanol and Hexane Solvents on Mortality of *Boophilus Microplus* Tick Larvae in Vitro. *JITV*, 10(02), 134-141.
- WHO (2005). *Guidelines for Laboratory and Field Testing of Mosquito Larvacides. World Health Organization Communicable Disease Control, Prevention, and Eradication Who Pesticide Evaluation Scheme*. Geneva: WHO Press, 1-41.
- Yudhastuti, R. & Vidiyani, A. (2005). Relationship of Environmental Conditions, Containers, and Community Behavior with the Existence of *Aedes Aegypti* Mosquito larvae in Endemic Areas of Dengue Hemorrhagic Fever in Surabaya, *Environmental Health Journal*, 1(2), 170-183.