



---

## PARENTAL KNOWLEDGE TOWARDS CHILDREN'S DENTAL AND ORAL HYGIENE IN SDLB 1 MARTAPURA

**Agno Elman Efzi ; Emilda Sari\***

*Department of Dental Hygiene, Poltekkes Kemenkes Banjarmasin ; Jl. Mistar Cokrokusumo No.1A ; Banjarbaru 70714 ; Indonesia*

### Abstract

Dental and oral hygiene problem is the leading cause of daily activities disruption in children such as missing school, bad learning concentration, appetite and nutritional intake. Knowledge of dental and oral hygiene should be given at early age in family by the parents. The aim of our work was to determine relationship between parents' knowledge and dental hygiene of the children. We undertook this research with analytical survey. A cross-sectional design was carried out. The sampling method was total sampling namely parents of children with special needs and the children as much as 80 people. Chi-Square test was performed to analyze the data. The results show that the knowledge of parents about dental and oral hygiene is not good (85%), while the dental and oral hygiene in children with special needs is poor (85%). Based on the Chi-Square Test, it produces a p-value of sig (2-sided) = 0.002 in the test, so it can obtain a p-value (probability value) from the test ( $p = 0.002 \leq 0.05$ ) so that  $H_0$  is rejected. In conclusion, a relationship between parental knowledge and oral hygiene in children with special needs does exist.

**Keywords:** *knowledge; dental and oral hygiene; children with special needs; parents*

### 1. Introduction

Caries is the most common chronic dental problem yet it is actually preventable (Syreen, et al. 2018; Alhabdan, et al., 2018). Caries attacked 3.5 billion people in the world and it is started when the first teeth appear at children age. The report of World Health Organization reveals 60 - 90% of school children worldwide has experienced caries especially in Asian countries

(World Health Organization, 2003). A study found that the caries prevalence and DMFT score of Indonesia are higher than other Southeast Asian countries (Zhang, et al. 2014; Jürgensen & Petersen, 2009; Urwannachotima & Hanvoravongchai, 2020; Lai, et al., 2018; Kubota, et al., 2017). It is proven that 61% of 12-year-old Indonesian children were affected by dental caries with the mean DMFT score in 1.58 (Maharani, et al., 2019).

Children may experience psychiatric and mental health issues. Special needs children are those under the age of 18 who have either

---

\*) Corresponding Author (Emilda Sari)

E-mail: [melda\\_ahg\\_bjm@yahoo.co.id](mailto:melda_ahg_bjm@yahoo.co.id)

physical or cognitive disabilities including intellectual disability (ID), Down Syndrome (DS), autism spectrum disorder (ASD), and attention deficit hyperactivity disorder (ADHD). Nevertheless, they still have the same dental problems of normal children such as dental caries and poor oral hygiene (Yeung, et al., 2019; Motaveloso et al., 2012; Vellappally et al., 2014). This condition can lead to dental aversion which associated with increase plaque levels and caries experience. Intellectual disabilities may also limit the self - care, resulting in poor dental hygiene (Sarvas, 2017). Children with special needs require more assistance because of their mental as well as physical challenges, even if they are over seven years old. Because some of them may learn slowly and being uncooperative, have difficulty to understand the behavior of dental hygiene such as brushing teeth (Ningrum, et al., 2021). It was mentioned that children with special need is prone having a high dental carries index and inadequate oral hygiene (Pini, et al., 2016).

The role of parents is essential in dental hygiene as they are the main caregivers to their children. The behavior of parents, particularly mothers can affect their children's health. Some of the behavior of the parents include tooth brushing habit, dietary, and food choices. A good behavior of the parents can affect the children in the effort of caries risk reduction (Bozorgmehr, et al., 2013). Family environment has something to do with the influence of good behavior about dental hygiene (Duijster, et al., 2015).

Therefore, the relationship of the parental knowledge regarding the dental hygiene of their children are very important to understand. Over the past ten years, the information of the relationship of the parent on the dental hygiene of the children have been conducted. A significant relationship of parental behavior and preschool children in Kerman, South East of Iran was proven in (Bozorgmehr et al., 2013). However, this study doesn't evaluate the knowledge of the parent among the relationship to the children dental hygiene, though it was mentioned that knowledge and attitude of the parents affect children dental health behavior and status. A survey on parental oral knowledge and behavior has been conducted in relationship to the children oral health in (Priya, et al., 2018; Abdat & Ramayana, 2020; Rogéria, et al., 2013). The results

indicate that the children poor oral health practice is reflected by the parent partial knowledge and behavior toward oral health. However, all aforementioned research suffer from a relationship to the children with special needs.

Therefore, our study conducted with 40 parents and their children in SDLB N (Sekolah Dasar Luar Biasa Negeri) 1 Martapura, Banjar Regency totaling 80 people. SDLB N 1 Martapura was established since 2018, located in Jl. Candra Kirana RT V/I, Martapura City is intended for children with special needs so that they can get basic services to help them getting access to education with different types of learning strategies and facilities.

Our objective was to evaluate the relationship of parental knowledge to the dental hygiene of the children with special needs. The study was performed by evaluating the knowledge of the parents by using questionnaire and the dental oral hygiene by examining OHI-S of the children. A further objective was exploring the parents' knowledge and the dental hygiene of the children with special needs.

## 2. Method

### Study design and sampling procedure

Our study type was analytical survey. The cross-sectional approach was used to determine the relationship between parental knowledge and oral hygiene and the approach and the variables of research were obtained in the same time. The study was conducted from October 2019 to May 2020. The variable of research consists of independent variable, i.e. knowledge of parents and dependent variable i.e. dental and oral hygiene of the children.

The population was all 40 students with special needs and 40 parents at SDLB N 1 Martapura, Banjar Regency. Total sampling was carried out with the total population, namely all students with special needs and their parents at SDLB 1 Martapura, Banjar Regency with a total sample of 80 people.

### Data collection

Primary data collection was carried out by conducting direct examinations of children with special needs at the SDLB N 1 Martapura, Banjar Regency using a diagnostic instrument set, OHI-S format and giving a dental and oral hygiene

knowledge questionnaire to parents of children with special needs. The questionnaire question was 19 in total. Each of question consists of 3 answers. Score 1 is given to the correct answer and score 0 is given to the incorrect answer. There are two categories in knowledge, i.e. Good if the score is 10 - 19, and poor if the score is 0 - 9.

Secondary data was data obtained from the research site, SDLB N 1 Martapura, Banjar Regency, regarding the names of students and their parents, age, gender, type of disability and the number of students with special needs and the parents.

### Data analysis

Data analysis was started by inputting the primary and secondary in frequency distribution table. After collecting the results of dental and oral hygiene examinations for children with special needs, then the results were statistically tested using the SPSS program with the Chi Square test to find out the existence of relationship in parental knowledge and oral hygiene (Mchugh, 2013). The category used for Debris Index and Calculus is based on in which good is scored 0 - 0.6; fair if scored 0.7 - 1.8; poor if 1.9 - 3.0 (Greene & Vermillion, 1964).

The conclusion of relationship was taken as follows: if the result of Chi Square test on p - value less than 0.05, meaning that there is a relationship between parental knowledge and dental and oral hygiene of the children with special need and vice versa, if p - value is more than 0.05, meaning that there is no relationship between parental knowledge and dental and oral hygiene of the children with special needs.

### 3. Result and Discussion

Results on frequency distribution are divided into distribution based on the parents and children. The data of parents include the number of respondents and knowledge. Meanwhile the data of children includes the number of respondents, and dental and oral hygiene.

**Table 1.** Frequency distribution of parent respondents

| Gender | Number of respondents | Percentage (%) |
|--------|-----------------------|----------------|
| Male   | 12                    | 30             |
| Female | 28                    | 70             |
| Total  | 40                    | 100            |

Table 1 shows the number of parents of children with special needs in which 30% of the respondents are male parent (father) and 70% are female parent (mother).

**Table 2.** Frequency distribution of students in SDLB N 1 Martapura

| Grade     | Number | Percentage (%) |
|-----------|--------|----------------|
| Grade I   | 12     | 30             |
| Grade II  | 10     | 25             |
| Grade III | 5      | 12.5           |
| Grade IV  | 4      | 10             |
| Grade V   | 4      | 10             |
| Grade VI  | 5      | 12.5           |
| Total     | 40     | 100            |

Table 2 shows the frequency distribution of the students that the total number of students in SDLB N 1 Martapura is 40.

**Table 3.** Frequency distribution of parental knowledge

| Category | Number of respondents | Percentage (%) |
|----------|-----------------------|----------------|
| Good     | 6                     | 15             |
| Poor     | 34                    | 85             |
| Total    | 40                    | 100            |

Table 3 states the number of parental knowledge with poor category is 34 people and good category is 6 people.

**Table 4.** Frequency distribution of dental and oral hygiene of the children

| Category | Number of Respondents | Percentage (%) |
|----------|-----------------------|----------------|
| Good     | 4                     | 10             |
| Fair     | 2                     | 5              |
| Poor     | 34                    | 85             |
| Total    | 40                    | 100            |

Table 4 shows the highest dental and oral hygiene percentage is in category of poor

with 34 children or 85% of the total respondents.

**Table 5.** Cross Tabulation of the Relationship between Parental Knowledge and Dental and Oral Hygiene in Children with Special Needs

| Parental knowledge | Dental and oral hygiene of the children |     |      |   |      |      | Total |     |
|--------------------|---|-----|------|---|------|------|-------|-----|
|                    | Good                                    |     | Fair |   | Poor |      | N     | %   |
|                    | N                                       | %   | N    | % | N    | %    |       |     |
| Good               | 1                                       | 2.5 | 2    | 5 | 3    | 7.5  | 6     | 15  |
| Poor               | 3                                       | 7.5 | 0    | 0 | 31   | 77.5 | 34    | 85  |
| Total              | 4                                       | 10  | 2    | 5 | 34   | 85   | 40    | 100 |

Table 5 shows from 40 parents there are 34 (85%) parents who have poor knowledge with children who have good dental and oral hygiene as many as 3 children (7.5%), and poor oral hygiene category as many as 31 children (77.5%). Meanwhile no children with fair category of dental and oral hygiene.

There are 6 parents (15%) who have good knowledge with children who have good dental and oral hygiene category as much as 1 child (2.5%), fair category as many as 2 children (5%) and the category of poor was 3 children (7.5%).

**Table 6.** Chi Square test results on parental knowledge and dental oral hygiene of the children

|                   | Value               | Df | Sig (2-sided) |
|-------------------|---------------------|----|---------------|
| Person Chi-Square | 12.664 <sup>a</sup> | 2  | 0.002         |

Table 6 shows the p value in the sig (2-sided) = 0.002 column in the test, meaning that the p value (probability value) of the test is less than, so H<sub>0</sub> is rejected. In conclusion, there is a relationship between parental knowledge and dental and oral hygiene in children with special needs at SDLB N 1 Martapura.

In table 5, the number of parents who have poor knowledge of dental and oral hygiene is higher than those who have good poor knowledge. As shown in the research of (Mahat & Bowen, 2017), parents actually have awareness that teeth are important to maintain. However

few understand the cavities appear in their children's teeth which cause cavities in permanent teeth. So, the bad dental and oral hygiene may be caused by baby teeth cavities that parents didn't realize.

One objective of Ministry of Health of Indonesia 2030 is that to reduce the proportion of dental caries experience of children at age 12 (Ministry of Health of Indonesia, 2019). WHO recommends three prevention of childhood caries i.e. primary prevention which includes primary health care programs especially for maternal and child health that is conducted in early age such as no sugars for baby until 2 years. Secondary prevention is performed focusing on early detection of carious lesions. This is focusing on the dental personnel and even mothers to detect early signs of carious lesions. Tertiary prevention includes reduce cavity by avoiding unnecessary extraction and restoration function. In tertiary prevention may apply rehabilitation for children if there is problem of child's behavior and cooperation (World Health Organization (WHO), 2016). This show that actually the starting point of caries prevention is by making sure that mother or the parents are having knowledge of dental and oral hygiene to promote good dental and oral hygiene to their own children. Since parental knowledge has significant influence on children's dental caries (Isong, et al., 2012).

Children with special healthcare needs (CSHCN) require health and related services more than that of children in general (McPherson et al., 1998). The children with special needs can suffer from mental and physical disabilities that influence them in mobility hindrance which resulting in higher poor dental hygiene (Bayarsaikhan, et al., 2015). This is in line with our study according to table 5 that either the parent have good knowledge, the highest category in dental and oral hygiene is poor. However, this is in accordance to the parental knowledge, which is in line with our study result in table 6 that there is relationship between parental knowledge and the dental and oral hygiene status of the children with special needs. Therefore, it is recommended that assessment of parental knowledge is important to perform first before the appropriate education program for the target is held (Mahat & Bowen,

2017). The poor dental and oral hygiene of the children with special needs is also caused by poor routine dentist visit, perception that dental care is expensive (Hendaus et al., 2020), socioeconomic (Oberoi, et al., 2016), educational background (Albatayneh, et al., 2019). The dental health personnel is needed to educate them, and it is recommended to initiate dental home process (Hendaus et al., 2020).

#### 4. Conclusion and Suggestion

We have presented a research to evaluate the relationship of parental knowledge and dental and oral hygiene status of the children with special needs in SDLB N 1 Martapura, Banjar Regency. The results comes into conclusion that there is a relationship between parental knowledge and dental and oral hygiene status of the children. It is shown by the results that the poor knowledge of the parent does in line with the poor dental and oral hygiene status of the children. We suggest that further research should be undertaken in the areas of factors related to the parental knowledge, so that the educational program held by dental personnel or Public Health Service is right on the target.

#### 5. Acknowledgments

We gratefully acknowledge the help provided by Poltekkes Kemenkes Banjarmasin, the Head of Education South Kalimantan Province, and SDLB N 1 Martapura, Banjar Regency for the cooperation and support.

#### 6. References

- Abdat, M., & Ramayana, I. (2020). Relationship between mother ' s knowledge and behaviour with oral health status of early childhood. *Padjadjaran Journal of Dentistry*, 32(3), 166-173. <https://doi.org/10.24198/pjd.vol32no2.24734>
- Al-batayneh, O. B., Al-khateeb, H. O., Ibrahim, W. M., & Khader, Y. S. (2019). Parental Knowledge and Acceptance of Different Treatment Options for Primary Teeth Provided by Dental Practitioners. *Frontiers in Public Health*, 7(November), 1-8. <https://doi.org/10.3389/fpubh.2019.00322>
- Alhabdan, Y. A., Albeshr, A. G., & Yenugadhathi, N. (2018). Prevalence of dental caries and associated factors among primary school children : a population-based cross-sectional study in Riyadh , Saudi Arabia. *Environmental Health and Preventive Medicine*, 23(60), 1-14.
- Bayarsaikhan, Z., Cruz, S., Neff, J., & Chi, D. L. (2015). Transitioning from pediatric care to adult care for adolescents with special health care needs: dentist perspectives (part 2). *Pediatr Dent.*, 37(5), 447-451.
- Bozorgmehr, E., Hajizamani, A., & Mohammadi, T. M. (2013). Oral Health Behavior of Parents as a Predictor of Oral Health Status of Their Children. *ISRN Dentistry*, (741783), 1-5. <https://doi.org/10.1155/2013/741783>
- Duijster, D., Jong-lenters, M. De, Verrips, E., & Loveren, C. Van. (2015). Establishing oral health promoting behaviours in children – parents ' views on barriers , facilitators and professional support : a qualitative study. *BMC Oral Health*, 15(157), 1-13. <https://doi.org/10.1186/s12903-015-0145-0>
- Greene, J. G., & Vermillion, J. R. (1964). The Simplified Oral Hygiene Index. *The Journal of The American Dental Association*, 68(1), 7-13. <https://doi.org/10.14219/jada.archive.1964.0034>
- Hendaus, M. A., Shaltout, D., Yasrab, D., Al-noubani, A., Hamad, S. G., Alamri, M., & Alhammadi, A. H. (2020). Parental Perception of a Dental Home for Children with Special Needs. *Pediatric Health, Medicine and Therapeutics*, 11, 379-384.
- Isong, I. A., Luff, D., Perrin, J. M., Winickoff, J. P., & Ng, M. W. (2012). Parental Perspectives of Early Childhood Caries. *Clinical Pediatrics*, 51(1), 77-85. <https://doi.org/10.1177/0009922811417856>
- Jürgensen, N., & Petersen, P. E. (2009). Oral health and the impact of socio-behavioural factors in a cross sectional survey of 12-year old school children in Laos. *BMC Oral Health*, 9(29), 1-11. <https://doi.org/10.1186/1472-6831-9-29>
- Kubota, Y., Jareonpituak, S., Narksawat, K., Satitvipawee, P., & Durward, C. (2017). Status , knowledge , attitudes , and practices of 12-18-year old orphaned children pertaining to dental caries in an dental clinic by non-governmental organization at Phnom. *Oral Health Care*, 2(2), 1-4. <https://doi.org/10.15761/OHC.1000117>
- Lai, B., Tan, W. K., & Lu, Q. S. (2018). Clinical efficacy of a two-year oral health

- programme for infants and toddlers in Singapore. *Singapore Med J*, 59(2), 87-93. <https://doi.org/10.11622/smedj.2017045>
- Maharani, D. A., Zhang, S., Gao, S. S., & Chu, C. (2019). Dental Caries and the Erosive Tooth Wear Status of 12-Year-Old Children in Jakarta , Indonesia. *International Journal of Environmental Research and Public Health*, 16(2994), 1-10. <https://doi.org/10.3390/ijerph16162994>
- Mahat, G., & Bowen, F. (2017). Parental Knowledge about Urban Preschool Children ' s Oral Health Risk. *Pediatric Nursing*, 43(1), 30-34.
- Mchugh, M. L. (2013). The Chi-square test of independence. *Biochemia Medica*, 23(2), 143-149. <https://doi.org/10.11613/BM.2013.018>
- McPherson, M., Arango, P., Fox, H., Lauver, C., McManus, M., Newacheck, P. W., ... Strickland, B. (1998). A New Definition of Children With Special Health Care Needs. *Pediatrics*, 102(1), 137-140.
- Ministry of Health of Indonesia. (2019). Faktor Risiko Kesehatan Gigi dan Mulut. *InfoDATIN*, 1-6. Retrieved from <https://pusdatin.kemkes.go.id/download.php?file=download/pusdatin/infodatin/infodatin-gigi.pdf>
- Mota-veloso, I., Nogueira, R., Eduardo, C., Alca, P., Aparecida, S., Ramos-jorge, M. L., & Oliveira-ferreira, F. (2012). Research in Developmental Disabilities Does intellectual disability affect the development of dental caries in patients with cerebral palsy ? *Research in Developmental Disabilities*, 33, 1503-1507. <https://doi.org/10.1016/j.ridd.2012.03.026>
- Ningrum, V., Bakar, A., Shieh, T., & Shih, Y. (2021). The Oral Health Inequities between Special Needs Children and Normal Children in Asia : A Systematic Review and. *Healthcare*, 9(410), 1-12. <https://doi.org/10.3390/healthcare9040410>
- Oberoi, S. S., Sharma, G., & Oberoi, A. (2016). A cross-sectional survey to assess the effect of socioeconomic status on the oral hygiene habits. *J Indian Soc Periodontol*, 20(5), 531-542. <https://doi.org/10.4103/0972-124X.201629>
- Pini, D. D. M., Fröhlich, P. C. G. R., & Rigo, L. (2016). Oral health evaluation in special needs individuals. *Einstein (Sao Paulo)*, 14(5554), 501-507. <https://doi.org/10.1590/S1679-45082016AO3712>
- Priya, M. M., Kumar, P., & Saraswathi, S. (2018). Assessment of parental oral health knowledge , attitude , and behavior. *Journal of Pharmacy Research*, 12(1), 10-13.
- Rogéria, A., Castilho, F. De, Luiz, F., Souza, T. De, & Puppim-rontani, R. M. (2013). Influence of family environment on children ' s oral health : a systematic review ☆. *Journal de Pediatria*, 89(2), 116-123. <https://doi.org/10.1016/j.jped.2013.03.014>
- Sarvas, E. (2017). Assessing Caries Risk in Children With Special Needs. *Decisions in Dentistry*, 3(3), 48-52.
- Syreen, S., Anwar, A., Ahmad, G., & Rahman, M. (2018). Prevalence of Dental Caries among Children Residing in Muslim Dominated. *Journal of Medical Science and Clinical Research*, 06(12), 817-819.
- Urwannachotima, N., & Hanvoravongchai, P. (2020). System dynamics analysis of dental caries status among Thai adults and elderly. *Journal of Health Research*, 34(2), 134-146. <https://doi.org/10.1108/JHR-05-2019-0097>
- Vellappally, S., Gardens, S. J., Kheraif, A. A. Al, Krishna, M., Babu, S., Hashem, M., ... Anil, S. (2014). The prevalence of malocclusion and its association with dental caries among 12-18-year-old disabled adolescents. *BMC Oral Health*, 14(123), 1-7.
- World Health Organization. (2003). *Continuous improvement of oral health in the 21st century – the approach of the WHO Global Oral Health Programme*. Geneva.
- World Health Organization (WHO). (2016). *WHO Expert Consultation on Public Health Intervention against Early Childhood Caries*. Retrieved from <https://apps.who.int/iris/bitstream/handle/10665/255627/WHO-NMH-PND-17.1-eng.pdf>
- Yeung, P. M., Wong, V. C. N., Mcgrath, C. P., Yiu, C. K. Y., & Lee, G. H. M. (2019). Oral health status of children with epilepsy in Hong Kong. *J Invest Clin Dent*, 10(February), 1-8. <https://doi.org/10.1111/jicd.12479>
- Zhang, S., Chau, A. M. H., Lo, E. C. M., & Chu, C. (2014). Dental caries and erosion status of 12-year-old Hong Kong children. *BMC Public Health*, 14(7), 1-7.



**JURNAL RISET KESEHATAN**

pISSN 2252-5068  
eISSN 2461-1026

Announcements | [read more >](#)

Call for Paper: Information for submission in Jurnal Riset Kesehatan in May 2022.

**Jurnal Riset Kesehatan**

JOURNAL HOME ISSUES INFORMATION OVERVIEW [SUBMIT](#) [SUBSCRIBE](#) search all articles

Home > User > Author > Submissions > #7625 > Summary

## #7625 Summary

SUMMARY REVIEW EDITING

### Submission

|                |  |
|----------------|--|
| Authors        | Agno Elman Efa, Emilda Sari  |
| Title          | PARENTAL KNOWLEDGE TOWARDS CHILDRENS DENTAL AND ORAL HYGIENE   |
| Original file  | 7625-22839-1-2M.DOCX 2021-09-07  |
| Supp. files    | 7625-22840-1-SP.PDF 2021-09-07<br>7625-22841-1-SP.PDF 2021-09-07<br>7625-22842-2-SP.PDF 2021-09-07<br>7625-22843-1-SP.PDF 2021-09-07 |
| Submitter      | Mrs. Emilda Sari   |
| Date submitted | September 7, 2021 - 01:12 PM   |
| Section        | Articles   |
| Editor         | M. Ibrahim Ayatullah<br>Diyah Fatmahan   |
| Abstract Views | 0  |



**PUBLISHER**  
Pusat Penelitian dan Pengabdian Kepada Masyarakat (PPPM)  
Poltekkes Kemenkes Semarang

- PEER REVIEWED
- AUTHOR GUIDELINES
- PUBLICATION ETHICS
- ONLINE SUBMISSION
- AUTHOR STATEMENT LETTER
- COPYRIGHT TRANSFER AGREEMENT

### Status

|               |  |
|---------------|--|
| Status        | Published - Vol 10, No 2 (2021): NOVEMBER 2021 |
| Initiated     | 2021-12-02                                     |
| Last modified | 2021-12-02                                     |

- COPYRIGHT TRANSFER AGREEMENT
- VIEW STATISTICS

### Submission Metadata

#### Authors

|   |                                |
|---|--------------------------------|
| Name  | Agno Elman Efa                 |
| Affiliation                                     | Poltekkes Kemenkes Banjarmasin |
| Country   | Indonesia                      |
| Bio Statement                                   | Department of Dental Hygiene   |
| Principal contact for editorial correspondence: |                                |
| Name  | Emilda Sari                    |
| Affiliation                                     | Poltekkes Kemenkes Banjarmasin |
| Country   | Indonesia                      |
| Bio Statement                                   | Department of Dental Hygiene   |

Select Language

English

[SUBMIT](#)

+62 899 1011 178

[jrk@poltekkes-smg.ac.id](mailto:jrk@poltekkes-smg.ac.id)

### Title and Abstract

|       |  |
|-------|--|
| Title | PARENTAL KNOWLEDGE TOWARDS CHILDRENS DENTAL AND ORAL HYGIENE |
|-------|--|

COLLABORATION WITH



Persatuan Perawat Nasional Indonesia

Abstract

Dental and oral hygiene problem is the leading cause of daily activities disruption in children such as missing school, bad learning concentration, appetite and nutritional intake. Knowledge of dental and oral hygiene should be given at early age in family by the parents. The aim of our work was to determine relationship between parents' knowledge and dental hygiene of the SDBL children (children with special needs). We undertook this research with analytical survey. A cross-sectional design was carried out. The sampling method was total sampling namely parents of children with special needs and the children as many as 80 people. Chi-Square test was performed to analyze the data. The results show that the knowledge of parents about dental and oral hygiene is not good (85%), while the dental and oral hygiene in children with special needs is poor (85%). Based on the Chi-Square Test, it produces a p-value of sig (2-sided) = 0.002 in the test, so it can obtain a p-value (probability value) from the test ( $p = 0.002 < \alpha = 0.05$ ) so that  $H_0$  is rejected. In conclusion, a relationship between parental knowledge and oral hygiene in children with special needs does exist.

Indexing

Keywords Knowledge; Dental and oral hygiene; Children with special needs; Parents  
Language en

Supporting Agencies

Agencies —



Perhimpunan Radiografer Indonesia



Persatuan Terapis Gigi dan Mulut Indonesia





Jurnal Riset Kesehatan, x (x), xxxx, 1 - 7

DOI: 10.31983/jrk.vvix.xxxx

**Jurnal Riset Kesehatan**

<http://ejournal.poltekkes-smg.ac.id/ojs/index.php/jrk>

## PARENTAL KNOWLEDGE TOWARDS CHILDREN'S DENTAL AND ORAL HYGIENE IN SDLB 1 MARTAPURA

**Agno Elman Efzi ; Emilda Sari\***

*Department of Dental Hygiene, Poltekkes Kemenkes Banjarmasin ; Jl. Mistar Cokrokusumo No.1A ; Banjarbaru 70714 ; Indonesia*

### Abstract

Dental and oral hygiene problem is the leading cause of daily activities disruption in children such as missing school, bad learning concentration, appetite and nutritional intake. Knowledge of dental and oral hygiene should be given at early age in family by the parents. The aim of our work was to determine relationship between parents' knowledge and dental hygiene of the children. We undertook this research with analytical survey. A cross-sectional design was carried out. The sampling method was total sampling namely parents of children with special needs and the children as much as 80 people. Chi-Square test was performed to analyze the data. The results show that the knowledge of parents about dental and oral hygiene is not good (85%), while the dental and oral hygiene in children with special needs is poor (85%). Based on the Chi-Square Test, it produces a p-value of sig (2-sided) = 0.002 in the test, so it can obtain a p-value (probability value) from the test ( $p = 0.002 < 0.05$ ) so that  $H_0$  is rejected. In conclusion, a relationship between parental knowledge and oral hygiene in children with special needs does exist.

**Keywords:** *knowledge; dental and oral hygiene; children with special needs; parents*

### 1. Introduction

Caries is the most common chronic dental problem yet it is actually preventable (Syreen, et al. 2018; Alhabdan, et al., 2018). Caries attacked 3.5 billion people in the world and it is started when the first teeth appear at children age. The report of World Health Organization reveals 60 – 90% of school children worldwide has experienced caries especially in Asian countries (World Health Organization, 2003). A study found that the caries prevalence and DMFT score

of Indonesia are higher than other Southeast Asian countries (Zhang, et al. 2014; Jürgensen & Petersen, 2009; Urwannachotima & Hanvoravongchai, 2020; Lai, et al., 2018; Kubota, et al., 2017). It is proven that 61% of 12-year-old Indonesian children were affected by dental caries with the mean DMFT score in 1.58 (Maharani, et al., 2019).

Children may experience psychiatric and mental health issues. Special needs children are those under the age of 18 who have either physical or cognitive disabilities including

intellectual disability (ID), Down Syndrome (DS), autism spectrum disorder (ASD), and attention deficit hyperactivity disorder (ADHD). Nevertheless, they still have the same dental problems of normal children such as dental caries and poor oral hygiene (Yeung, et al., 2019; Motaveloso et al., 2012; Vellappally et al., 2014). This condition can lead to dental aversion which associated with increase plaque levels and caries experience. Intellectual disabilities may also limit the self - care, resulting in poor dental hygiene (Sarvas, 2017). Children with special needs require more assistance because of their mental as well as physical challenges, even if they are over seven years old. Because some of them may learn slowly and being uncooperative, have difficulty to understand the behavior of dental hygiene such as brushing teeth (Ningrum, et al., 2021). It was mentioned that children with special need is prone having a high dental carries index and inadequate oral hygiene (Pini, et al., 2016).

The role of parents is essential in dental hygiene as they are the main caregivers to their children. The behavior of parents, particularly mothers can affect their children's health. Some of the behavior of the parents include tooth brushing habit, dietary, and food choices. A good behavior of the parents can affect the children in the effort of caries risk reduction (Bozorgmehr, et al., 2013). Family environment has something to do with the influence of good behavior about dental hygiene (Duijster, et al., 2015).

Therefore, the relationship of the parental knowledge regarding the dental hygiene of their children are very important to understand. Over the past ten years, the information of the relationship of the parent on the dental hygiene of the children have been conducted. A significant relationship of parental behavior and preschool children in Kerman, South East of Iran was proven in (Bozorgmehr et al., 2013). However, this study doesn't evaluate the knowledge of the parent among the relationship to the children dental hygiene, though it was mentioned that knowledge and attitude of the parents affect children dental health behavior and status. A survey on parental oral knowledge and behavior has been conducted in relationship to the children oral health in (Priya, et al., 2018; Abdat & Ramayana, 2020; Rogéria, et al., 2013). The results indicate that the children poor oral health practice

is reflected by the parent partial knowledge and behavior toward oral health. However, all aforementioned research suffer from a relationship to the children with special needs.

Therefore, our study conducted with 40 parents and their children in SDLB N (Sekolah Dasar Luar Biasa Negeri) 1 Martapura, Banjar Regency totaling 80 people. SDLB N 1 Martapura was established since 2018, located in Jl. Candra Kirana RT V/I, Martapura City is intended for children with special needs so that they can get basic services to help them getting access to education with different types of learning strategies and facilities.

Our objective was to evaluate the relationship of parental knowledge to the dental hygiene of the children with special needs. The study was performed by evaluating the knowledge of the parents by using questionnaire and the dental oral hygiene by examining OHI-S of the children. A further objective was exploring the parents' knowledge and the dental hygiene of the children with special needs.

## 2. Method

### Study design and sampling procedure

Our study type was analytical survey. The cross-sectional approach was used to determine the relationship between parental knowledge and oral hygiene and the approach and the variables of research were obtained in the same time. The study was conducted from October 2019 to May 2020. The variable of research consists of independent variable, i.e. knowledge of parents and dependent variable i.e. dental and oral hygiene of the children.

The population was all 40 students with special needs and 40 parents at SDLB N 1 Martapura, Banjar Regency. Total sampling was carried out with the total population, namely all students with special needs and their parents at SDLB 1 Martapura, Banjar Regency with a total sample of 80 people.

### Data collection

Primary data collection was carried out by conducting direct examinations of children with special needs at the SDLB N 1 Martapura, Banjar Regency using a diagnostic instrument set, OHI-S format and giving a dental and oral hygiene knowledge questionnaire to parents of children

with special needs. The questionnaire question was 19 in total. Each of question consists of 3 answers. Score 1 is given to the correct answer and score 0 is given to the incorrect answer. There are two categories in knowledge, i.e. Good if the score is 10 - 19, and poor if the score is 0 - 9.

Secondary data was data obtained from the research site, SDLB N 1 Martapura, Banjar Regency, regarding the names of students and their parents, age, gender, type of disability and the number of students with special needs and the parents.

**Data analysis**

Data analysis was started by inputting the primary and secondary in frequency distribution table. After collecting the results of dental and oral hygiene examinations for children with special needs, then the results were statistically tested using the SPSS program with the Chi Square test to find out the existence of relationship in parental knowledge and oral hygiene (Mchugh, 2013). The category used for Debris Index and Calculus is based on in which good is scored 0 - 0.6; fair if scored 0.7 - 1.8; poor if 1.9 - 3.0 (Greene & Vermillion, 1964).

The conclusion of relationship was taken as follows: if the result of Chi Square test on p - value less than 0.05, meaning that there is a relationship between parental knowledge and dental and oral hygiene of the children with special need and vice versa, if p - value is more than 0.05, meaning that there is no relationship between parental knowledge and dental and oral hygiene of the children with special needs.

**3. Result and Discussion**

Results on frequency distribution are divided into distribution based on the parents and children. The data of parents include the number of respondents and knowledge. Meanwhile the data of children includes the number of respondents, and dental and oral hygiene.

**Table 1.** Frequency distribution of parent respondents

| Gender | Number of respondents | Percentage (%) |
|--------|-----------------------|----------------|
| Male   | 12                    | 30             |
| Female | 28                    | 70             |
| Total  | 40                    | 100            |

Table 1 shows the number of parents of children with special needs in which 30% of the respondents are male parent (father) and 70% are female parent (mother).

**Table 2.** Frequency distribution of students in SDLB N 1 Martapura

| Grade     | Number | Percentage (%) |
|-----------|--------|----------------|
| Grade I   | 12     | 30             |
| Grade II  | 10     | 25             |
| Grade III | 5      | 12.5           |
| Grade IV  | 4      | 10             |
| Grade V   | 4      | 10             |
| Grade VI  | 5      | 12.5           |
| Total     | 40     | 100            |

Table 2 shows the frequency distribution of the students that the total number of students in SDLB N 1 Martapura is 40.

**Table 3.** Frequency distribution of parental knowledge

| Category | Number of respondents | Percentage (%) |
|----------|-----------------------|----------------|
| Good     | 6                     | 15             |
| Poor     | 34                    | 85             |
| Total    | 40                    | 100            |

Table 3 states the number of parental knowledge with poor category is 34 people and good category is 6 people.

**Table 4.** Frequency distribution of dental and oral hygiene of the children

| Category | Number of Respondents | Percentage (%) |
|----------|-----------------------|----------------|
| Good     | 4                     | 10             |
| Fair     | 2                     | 5              |
| Poor     | 34                    | 85             |
| Total    | 40                    | 100            |

Table 4 shows the highest dental and oral hygiene percentage is in category of poor with 34 children or 85% of the total respondents.

**Table 5.** Cross Tabulation of the Relationship between Parental Knowledge and Dental and Oral Hygiene in Children with Special Needs

| Parental knowledge | Dental and oral hygiene of the children |     |      |   |      |      | Total |     |
|--------------------|---|-----|------|---|------|------|-------|-----|
|                    | Good                                    |     | Fair |   | Poor |      | N     | %   |
|                    | N                                       | %   | N    | % | N    | %    |       |     |
| Good               | 1                                       | 2.5 | 2    | 5 | 3    | 7.5  | 6     | 15  |
| Poor               | 3                                       | 7.5 | 0    | 0 | 31   | 77.5 | 34    | 85  |
| Total              | 4                                       | 10  | 2    | 5 | 34   | 85   | 40    | 100 |

Table 5 shows from 40 parents there are 34 (85%) parents who have poor knowledge with children who have good dental and oral hygiene as many as 3 children (7.5%), and poor oral hygiene category as many as 31 children (77.5%). Meanwhile no children with fair category of dental and oral hygiene.

There are 6 parents (15%) who have good knowledge with children who have good dental and oral hygiene category as much as 1 child (2.5%), fair category as many as 2 children (5%) and the category of poor was 3 children (7.5%).

**Table 6.** Chi Square test results on parental knowledge and dental oral hygiene of the children

|                   | Value               | Df | Sig (2-sided) |
|-------------------|---------------------|----|---------------|
| Person Chi-Square | 12.664 <sup>a</sup> | 2  | 0.002         |

Table 6 shows the p value in the sig (2-sided) = 0.002 column in the test, meaning that the p value (probability value) of the test is less than, so H<sub>0</sub> is rejected. In conclusion, there is a relationship between parental knowledge and dental and oral hygiene in children with special needs at SDLB N 1 Martapura.

In table 5, the number of parents who have poor knowledge of dental and oral hygiene is higher than those who have good poor knowledge. As shown in the research of (Mahat & Bowen, 2017), parents actually have awareness that teeth are important to maintain. However few understand the cavities appear in their children's teeth which cause cavities in permanent teeth. So, the bad dental and oral

hygiene may be caused by baby teeth cavities that parents didn't realize.

One objective of Ministry of Health of Indonesia 2030 is that to reduce the proportion of dental caries experience of children at age 12 (Ministry of Health of Indonesia, 2019). WHO recommends three prevention of childhood caries i.e. primary prevention which includes primary health care programs especially for maternal and child health that is conducted in early age such as no sugars for baby until 2 years. Secondary prevention is performed focusing on early detection of carious lesions. This is focusing on the dental personnel and even mothers to detect early signs of carious lesions. Tertiary prevention includes reduce cavity by avoiding unnecessary extraction and restoration function. In tertiary prevention may apply rehabilitation for children if there is problem of child's behavior and cooperation (World Health Organization (WHO), 2016). This show that actually the starting point of caries prevention is by making sure that mother or the parents are having knowledge of dental and oral hygiene to promote good dental and oral hygiene to their own children. Since parental knowledge has significant influence on children's dental caries (Isong, et al., 2012).

Children with special healthcare needs (CSHCN) require health and related services more than that of children in general (McPherson et al., 1998). The children with special needs can suffer from mental and physical disabilities that influence them in mobility hindrance which resulting in higher poor dental hygiene (Bayarsaikhan, et al., 2015). This is in line with our study according to table 5 that either the parent have good knowledge, the highest category in dental and oral hygiene is poor. However, this is in accordance to the parental knowledge, which is in line with our study result in table 6 that there is relationship between parental knowledge and the dental and oral hygiene status of the children with special needs. Therefore, it is recommended that assessment of parental knowledge is important to perform first before the appropriate education program for the target is held (Mahat & Bowen, 2017). The poor dental and oral hygiene of the children with special needs is also caused by poor routine dentist visit, perception that dental care is

**Commented [r1]:** Pengetahuan & OHIS blh dibuat satu tabel sehingga dpt terlihat p value ,chi square Dan juga OR

Jadi bisa disebutkan jika seorang berpengetahuan baik akan lbh meningkat derajat kebersihan mulut nya

expensive (Hendaus et al., 2020), socioeconomic (Oberoi, et al., 2016), educational background (Albatayneh, et al., 2019). The dental health personnel is needed to educate them, and it is recommended to initiate dental home process (Hendaus et al., 2020).

#### 4. Conclusion and Suggestion

We have presented a research to evaluate the relationship of parental knowledge and dental and oral hygiene status of the children with special needs in SDLB N 1 Martapura, Banjar Regency. The results comes into conclusion that there is a relationship between parental knowledge and dental and oral hygiene status of the children. It is shown by the results that the poor knowledge of the parent does in line with the poor dental and oral hygiene status of the children. We suggest that further research should be undertaken in the areas of factors related to the parental knowledge, so that the educational program held by dental personnel or Public Health Service is right on the target.

#### 5. Acknowledgments

We gratefully acknowledge the help provided by Poltekkes Kemenkes Banjarmasin, the Head of Education South Kalimantan Province, and SDLB N 1 Martapura, Banjar Regency for the cooperation and support.

#### 6. References

- Abdat, M., & Ramayana, I. (2020). Relationship between mother ' s knowledge and behaviour with oral health status of early childhood. *Padjadjaran Journal of Dentistry*, 32(3), 166-173. <https://doi.org/10.24198/pjd.vol32no2.24734>
- Al-batayneh, O. B., Al-khateeb, H. O., Ibrahim, W. M., & Khader, Y. S. (2019). Parental Knowledge and Acceptance of Different Treatment Options for Primary Teeth Provided by Dental Practitioners. *Frontiers in Public Health*, 7(November), 1-8. <https://doi.org/10.3389/fpubh.2019.00322>
- Alhabdan, Y. A., Albeshr, A. G., & Yenugadhathi, N. (2018). Prevalence of dental caries and associated factors among primary school children : a population-based cross-sectional study in Riyadh , Saudi Arabia. *Environmental Health and Preventive Medicine*, 23(60), 1-14.
- Bayarsaikhan, Z., Cruz, S., Neff, J., & Chi, D. L. (2015). Transitioning from pediatric care to adult care for adolescents with special health care needs: dentist perspectives (part 2). *Pediatr Dent.*, 37(5), 447-451.
- Bozorgmehr, E., Hajizamani, A., & Mohammadi, T. M. (2013). Oral Health Behavior of Parents as a Predictor of Oral Health Status of Their Children. *ISRN Dentistry*, (741783), 1-5. <https://doi.org/10.1155/2013/741783>
- Duijster, D., Jong-lenters, M. De, Verrips, E., & Loveren, C. Van. (2015). Establishing oral health promoting behaviours in children - parents ' views on barriers , facilitators and professional support : a qualitative study. *BMC Oral Health*, 15(157), 1-13. <https://doi.org/10.1186/s12903-015-0145-0>
- Greene, J. G., & Vermillion, J. R. (1964). The Simplified Oral Hygiene Index. *The Journal of The American Dental Association*, 68(1), 7-13. <https://doi.org/10.14219/jada.archive.1964.0034>
- Hendaus, M. A., Shaltout, D., Yasrab, D., Al-noubani, A., Hamad, S. G., Alamri, M., & Alhammadi, A. H. (2020). Parental Perception of a Dental Home for Children with Special Needs. *Pediatric Health, Medicine and Therapeutics*, 11, 379-384.
- Isong, I. A., Luff, D., Perrin, J. M., Winickoff, J. P., & Ng, M. W. (2012). Parental Perspectives of Early Childhood Caries. *Clinical Pediatrics*, 51(1), 77-85. <https://doi.org/10.1177/0009922811417856>
- Jürgensen, N., & Petersen, P. E. (2009). Oral health and the impact of socio-behavioural factors in a cross sectional survey of 12-year old school children in Laos. *BMC Oral Health*, 9(29), 1-11. <https://doi.org/10.1186/1472-6831-9-29>
- Kubota, Y., Jareonpituak, S., Narksawat, K., Satitvipawee, P., & Durward, C. (2017). Status , knowledge , attitudes , and practices of 12-18-year old orphaned children pertaining to dental caries in an dental clinic by non-governmental organization at Phnom. *Oral Health Care*, 2(2), 1-4. <https://doi.org/10.15761/OHC.1000117>
- Lai, B., Tan, W. K., & Lu, Q. S. (2018). Clinical efficacy of a two-year oral health programme for infants and toddlers in Singapore. *Singapore Med J*, 59(2), 87-93. <https://doi.org/10.11622/smedj.2017045>

**Commented [r2]:** Saran yg terbaik buat org tua

Kesimpulan  
Value dari Penelitian ini

- Maharani, D. A., Zhang, S., Gao, S. S., & Chu, C. (2019). Dental Caries and the Erosive Tooth Wear Status of 12-Year-Old Children in Jakarta, Indonesia. *International Journal of Environmental Research and Public Health*, 16(2994), 1-10. <https://doi.org/10.3390/ijerph16162994>
- Mahat, G., & Bowen, F. (2017). Parental Knowledge about Urban Preschool Children's Oral Health Risk. *Pediatric Nursing*, 43(1), 30-34.
- Mchugh, M. L. (2013). The Chi-square test of independence. *Biochemia Medica*, 23(2), 143-149. <https://doi.org/10.11613/BM.2013.018>
- McPherson, M., Arango, P., Fox, H., Lauver, C., McManus, M., Newacheck, P. W., ... Strickland, B. (1998). A New Definition of Children With Special Health Care Needs. *Pediatrics*, 102(1), 137-140.
- Ministry of Health of Indonesia. (2019). Faktor Risiko Kesehatan Gigi dan Mulut. *InfoDATIN*, 1-6. Retrieved from <https://pusdatin.kemkes.go.id/download.php?file=download/pusdatin/infodatin/infodatin-gigi.pdf>
- Mota-veloso, I., Nogueira, R., Eduardo, C., Alca, P., Aparecida, S., Ramos-jorge, M. L., & Oliveira-ferreira, F. (2012). Research in Developmental Disabilities Does intellectual disability affect the development of dental caries in patients with cerebral palsy? *Research in Developmental Disabilities*, 33, 1503-1507. <https://doi.org/10.1016/j.ridd.2012.03.026>
- Ningrum, V., Bakar, A., Shieh, T., & Shih, Y. (2021). The Oral Health Inequities between Special Needs Children and Normal Children in Asia: A Systematic Review and. *Healthcare*, 9(410), 1-12. <https://doi.org/10.3390/healthcare9040410>
- Oberoi, S. S., Sharma, G., & Oberoi, A. (2016). A cross-sectional survey to assess the effect of socioeconomic status on the oral hygiene habits. *J Indian Soc Periodontol*, 20(5), 531-542. <https://doi.org/10.4103/0972-124X.201629>
- Pini, D. D. M., Fröhlich, P. C. G. R., & Rigo, L. (2016). Oral health evaluation in special needs individuals. *Einstein (Sao Paulo)*, 14(55), 501-507. <https://doi.org/10.1590/S1679-45082016AO3712>
- Priya, M. M., Kumar, P., & Saraswathi, S. (2018). Assessment of parental oral health knowledge, attitude, and behavior. *Journal of Pharmacy Research*, 12(1), 10-13.
- Rogéria, A., Castilho, F. De, Luiz, F., Souza, T. De, & Puppim-rontani, R. M. (2013). Influence of family environment on children's oral health: a systematic review. *Journal de Pediatria*, 89(2), 116-123. <https://doi.org/10.1016/j.jped.2013.03.014>
- Sarvas, E. (2017). Assessing Caries Risk in Children With Special Needs. *Decisions in Dentistry*, 3(3), 48-52.
- Syreen, S., Anwar, A., Ahmad, G., & Rahman, M. (2018). Prevalence of Dental Caries among Children Residing in Muslim Dominated. *Journal of Medical Science and Clinical Research*, 06(12), 817-819.
- Urwannachotima, N., & Hanvoravongchai, P. (2020). System dynamics analysis of dental caries status among Thai adults and elderly. *Journal of Health Research*, 34(2), 134-146. <https://doi.org/10.1108/JHR-05-2019-0097>
- Vellappally, S., Gardens, S. J., Kheraif, A. A. Al, Krishna, M., Babu, S., Hashem, M., ... Anil, S. (2014). The prevalence of malocclusion and its association with dental caries among 12-18-year-old disabled adolescents. *BMC Oral Health*, 14(123), 1-7.
- World Health Organization. (2003). *Continuous improvement of oral health in the 21st century - the approach of the WHO Global Oral Health Programme*. Geneva.
- World Health Organization (WHO). (2016). *WHO Expert Consultation on Public Health Intervention against Early Childhood Caries*. Retrieved from <https://apps.who.int/iris/bitstream/handle/10665/255627/WHO-NMH-PND-17.1-eng.pdf>
- Yeung, P. M., Wong, V. C. N., Mcgrath, C. P., Yiu, C. K. Y., & Lee, G. H. M. (2019). Oral health status of children with epilepsy in Hong Kong. *J Invest Clin Dent*, 10(February), 1-8. <https://doi.org/10.1111/jicd.12479>
- Zhang, S., Chau, A. M. H., Lo, E. C. M., & Chu, C. (2014). Dental caries and erosion status of 12-year-old Hong Kong children. *BMC Public Health*, 14(7), 1-7.

Catatan editor:

1. Masukan dari editorial team dan reviewer untk judul menghapus lokasi SDLB 1 Martapura, sehingga judul menjadi Parental Knowledge Towards Children's Dental and Oral Hygiene.
2. Hasil final editorial untuk menghapus hasil tabel kuesioner dan hasil uji validity. Hal tsb masuk pada supplementary file saja
3. Untuk tabel frekuensi dirasa cukup dengan deskripsi saja. Cukup dengan tabel cross tab dan chisquare
4. Tabel 4 tidak perlu dirinci grade nya sudah tertuang dalam metode bahwa populasi adalah seluruh SD dengan jumlah 40



## PARENTAL KNOWLEDGE TOWARDS CHILDREN'S DENTAL AND ORAL HYGIENE IN SDLB 1 MARTAPURA

Agno Elman Efzi ; Emilda Sari\*

Department of Dental Hygiene, Poltekkes Kemenkes Banjarmasin ; Jl. Mistar Cokrokusumo No.1A ; Banjarbaru 70714 ; Indonesia

### Abstract

Dental and oral hygiene problem is the leading cause of daily activities disruption in children such as missing school, bad learning concentration, appetite and nutritional intake. Knowledge of dental and oral hygiene should be given at early age in family by the parents. The aim of our work was to determine relationship between parents' knowledge and dental hygiene of the children. We undertook this research with analytical survey. A cross-sectional design was carried out. The sampling method was total sampling namely parents of children with special needs and the children as much as 80 people. Chi-Square test was performed to analyze the data. The results show that the knowledge of parents about dental and oral hygiene is not good (85%), while the dental and oral hygiene in children with special needs is poor (85%). Based on the Chi-Square Test, it produces a p-value of sig (2-sided) = 0.002 in the test, so it can obtain a p-value (probability value) from the test ( $p = 0.002 < = 0.05$ ) so that  $H_0$  is rejected. In conclusion, a relationship between parental knowledge and oral hygiene in children with special needs does exist.

**Keywords:** *knowledge; dental and oral hygiene; children with special needs; parents*

### 1. Introduction

Caries is the most common chronic dental problem yet it is actually preventable (Syreen, et al. 2018; Alhabdan, et al., 2018). Caries attacked 3.5 billion people in the world and it is started when the first teeth appear at children age. The report of World Health Organization reveals 60 - 90% of school children worldwide has experienced caries especially in Asian countries (World Health Organization, 2003). A study found that the caries prevalence and DMFT score of Indonesia are higher than other Southeast Asian countries (Zhang, et al. 2014; Jürgensen &

Petersen, 2009; Urwannachotima & Hanvoravongchai, 2020; Lai, et al., 2018; Kubota, et al., 2017). It is proven that 61% of 12-year-old Indonesian children were affected by dental caries with the mean DMFT score in 1.58 (Maharani, et al., 2019).

Children may experience psychiatric and mental health issues. Special needs children are those under the age of 18 who have either physical or cognitive disabilities including intellectual disability (ID), Down Syndrome (DS), autism spectrum disorder (ASD), and attention deficit hyperactivity disorder (ADHD).

**Commented [a1]:** Why not mention about SLB students...not general students

**Commented [a2]:** Countable...

**Commented [a3]:** Explain...this prevalence in what group..children or adult or...?



Nevertheless, they still have the same dental problems of normal children such as dental caries and poor oral hygiene (Yeung, et al., 2019; Motaveloso et al., 2012; Vellappally et al., 2014). This condition can lead to dental aversion which associated with increase plaque levels and caries experience. Intellectual disabilities may also limit the self - care, resulting in poor dental hygiene (Sarvas, 2017). Children with special needs require more assistance because of their mental as well as physical challenges, even if they are over seven years old. Because some of them may learn slowly and being uncooperative, have difficulty to understand the behavior of dental hygiene such as brushing teeth (Ningrum, et al., 2021). It was mentioned that children with special need is prone having a high dental carries index and inadequate oral hygiene (Pini, et al., 2016).

The role of parents is essential in dental hygiene as they are the main caregivers to their children. The behavior of parents, particularly mothers can affect their children's health. Some of the behavior of the parents include tooth brushing habit, dietary, and food choices. A good behavior of the parents can affect the children in the effort of caries risk reduction(Bozorgmehr, et al., 2013). Family environment has something to do with the influence of good behavior about dental hygiene(Duijster, et al., 2015).

Therefore, the relationship of the parental knowledge regarding the dental hygiene of their children are very important to understand. Over the past ten years, the information of the relationship of the parent on the dental hygiene of the children have been conducted. A significant relationship of parental behavior and preschool children in Kerman, South East of Iran was proven in (Bozorgmehr et al., 2013). However, this study doesn't evaluate the knowledge of the parent among the relationship to the children dental hygiene, though it was mentioned that knowledge and attitude of the parents affect children dental health behavior and status. A survey on parental oral knowledge and behavior has been conducted in relationship to the children oral health in (Priya, et al., 2018; Abdat & Ramayana, 2020; Rogéria, et al., 2013). The results indicate that the children poor oral health practice is reflected by the parent partial knowledge and behavior toward oral health. However, all

aforementioned research suffer from a relationship to the children with special needs.

Therefore, our study conducted with 40 parents and their children in SDLB N (Sekolah Dasar Luar Biasa Negeri) 1 Martapura, Banjar Regency totaling 80 people. SDLB N 1 Martapura was established since 2018, located in Jl. Candra Kirana RT V/I, Martapura City is intended for children with special needs so that they can get basic services to help them getting access to education with different types of learning strategies and facilities.

Our objective was to evaluate the relationship of parental knowledge to the dental hygiene of the children with special needs. The study was performed by evaluating the knowledge of the parents by using questionnaire and the dental oral hygiene by examining OHI-S of the children. A further objective was exploring the parents' knowledge and the dental hygiene of the children with special needs.

## 2. Method

### Study design and sampling procedure

Our study type was analytical survey. The cross-sectional approach was used to determine the relationship between parental knowledge and oral hygiene and the approach and the variables of research were obtained in the same time. The study was conducted from October 2019 to May 2020. The variable of research consists of independent variable, i.e. knowledge of parents and dependent variable i.e. dental and oral hygiene of the children.

The population was all 40 students with special needs and 40 parents at SDLB N 1 Martapura, Banjar Regency. Total sampling was carried out with the total population, namely all students with special needs and their parents at SDLB 1 Martapura, Banjar Regency with a total sample of 80 people.

### Data collection

Primary data collection was carried out by conducting direct examinations of children with special needs at the SDLB N 1 Martapura, Banjar Regency using a diagnostic instrument set, OHI-S format and giving a dental and oral hygiene knowledge questionnaire to parents of children with special needs. The questionnaire question was 19 in total. Each of question consists of 3

**Commented [a4]:** The same or higher risk...?? as their condition which need support from others

**Commented [a6]:** Why with therefore..??

**Commented [a7]:** It is method...not suppose write at background

**Commented [a8]:** What is the purpose of this sentence..?? explore..??

**Commented [a5]:** Prep...not common in early sentence

**Commented [a9]:** Mention the contain of questionnaire of knowledge

**Commented [a10]:** The validity of quest..??contain question??

answers. Score 1 is given to the correct answer and score 0 is given to the incorrect answer. There are two categories in knowledge, i.e. Good if the score is 10 – 19, and poor if the score is 0 – 9.

Secondary data was data obtained from the research site, SDLB N 1 Martapura, Banjar Regency, regarding the names of students and their parents, age, gender, type of disability and the number of students with special needs and the parents.

**Data analysis**

Data analysis was started by inputting the primary and secondary in frequency distribution table. After collecting the results of dental and oral hygiene examinations for children with special needs, then the results were statistically tested using the SPSS program with the Chi Square test to find out the existence of relationship in parental knowledge and oral hygiene (Mchugh, 2013). The category used for Debris Index and Calculus is based on in which good is scored 0 – 0.6; fair if scored 0.7 – 1.8; poor if 1.9 – 3.0 (Greene & Vermillion, 1964).

The conclusion of relationship was taken as follows: if the result of Chi Square test on p - value less than 0.05, meaning that there is a relationship between parental knowledge and dental and oral hygiene of the children with special need and vice versa, if p - value is more than 0.05, meaning that there is no relationship between parental knowledge and dental and oral hygiene of the children with special needs.

**3. Result and Discussion**

Results on frequency distribution are divided into distribution based on the parents and children. The data of parents include the number of respondents and knowledge. Meanwhile the data of children includes the number of respondents, and dental and oral hygiene.

**Table 1.** Frequency distribution of parent respondents

| Gender | Number of respondents | Percentage (%) |
|--------|-----------------------|----------------|
| Male   | 12                    | 30             |

|        |    |     |
|--------|----|-----|
| Female | 28 | 70  |
| Total  | 40 | 100 |

Table 1 shows the number of parents of children with special needs in which 30% of the respondents are male parent (father) and 70% are female parent (mother).

**Table 2.** Frequency distribution of students in SDLB N 1 Martapura

| Grade     | Number | Percentage (%) |
|-----------|--------|----------------|
| Grade I   | 12     | 30             |
| Grade II  | 10     | 25             |
| Grade III | 5      | 12.5           |
| Grade IV  | 4      | 10             |
| Grade V   | 4      | 10             |
| Grade VI  | 5      | 12.5           |
| Total     | 40     | 100            |

Table 2 shows the frequency distribution of the students that the total number of students in SDLB N 1 Martapura is 40.

**Table 3.** Frequency distribution of parental knowledge

| Category | Number of respondents | Percentage (%) |
|----------|-----------------------|----------------|
| Good     | 6                     | 15             |
| Poor     | 34                    | 85             |
| Total    | 40                    | 100            |

Table 3 states the number of parental knowledge with poor category is 34 people and good category is 6 people.

**Table 4.** Frequency distribution of dental and oral hygiene of the children

| Category | Number of Respondents | Percentage (%) |
|----------|-----------------------|----------------|
| Good     | 4                     | 10             |
| Fair     | 2                     | 5              |
| Poor     | 34                    | 85             |
| Total    | 40                    | 100            |

Table 4 shows the highest dental and oral hygiene percentage is in category of poor with 34 children or 85% of the total respondents.

**Table 5.** Cross Tabulation of the Relationship between Parental Knowledge and Dental and

**Commented [a13]:** Why not all mother who look after more the children with special need??

**Commented [a11]:** Not necessary ..this sentence

**Commented [a12]:** All tables should be follow rules of tables for publication

Oral Hygiene in Children with Special Needs

| Parental knowledge | Dental and oral hygiene of the children |     |      |   |      |      | Total |     |
|--------------------|---|-----|------|---|------|------|-------|-----|
|                    | Good                                    |     | Fair |   | Poor |      | n     | %   |
|                    | n                                       | %   | n    | % | n    | %    |       |     |
| Good               | 1                                       | 2.5 | 2    | 5 | 3    | 7.5  | 6     | 15  |
| Poor               | 3                                       | 7.5 | 0    | 0 | 31   | 77.5 | 34    | 85  |
| Total              | 4                                       | 10  | 2    | 5 | 34   | 85   | 40    | 100 |

Table 5 shows from 40 parents there are 34 (85%) parents who have poor knowledge with children who have good dental and oral hygiene as many as 3 children (7.5%), and poor oral hygiene category as many as 31 children (77.5%). Meanwhile no children with fair category of dental and oral hygiene.

There are 6 parents (15%) who have good knowledge with children who have good dental and oral hygiene category as much as 1 child (2.5%), fair category as many as 2 children (5%) and the category of poor was 3 children (7.5%).

**Table 6.** Chi Square test results on parental knowledge and dental oral hygiene of the children

|                   | Value               | Df | Sig (2-sided) |
|-------------------|---------------------|----|---------------|
| Person Chi-Square | 12.664 <sup>a</sup> | 2  | 0.002         |

Table 6 shows the p value in the sig (2-sided) = 0.002 column in the test, meaning that the p value (probability value) of the test is less than, so H<sub>0</sub> is rejected. In conclusion, there is a relationship between parental knowledge and dental and oral hygiene in children with special needs at SDLB N 1 Martapura.

In table 5, the number of parents who have poor knowledge of dental and oral hygiene is higher than those who have good poor knowledge. As shown in the research of (Mahat & Bowen, 2017), parents actually have awareness that teeth are important to maintain. However few understand the cavities appear in their children's teeth which cause cavities in permanent teeth. So, the bad dental and oral hygiene may be caused by baby teeth cavities that parents didn't realize.

One objective of Ministry of Health of Indonesia 2030 is that to reduce the proportion of dental caries experience of children at age 12 (Ministry of Health of Indonesia, 2019). WHO recommends three prevention of childhood caries i.e. primary prevention which includes primary health care programs especially for maternal and child health that is conducted in early age such as no sugars for baby until 2 years. Secondary prevention is performed focusing on early detection of carious lesions. This is focusing on the dental personnel and even mothers to detect early signs of carious lesions. Tertiary prevention includes reduce cavity by avoiding unnecessary extraction and restoration function. In tertiary prevention may apply rehabilitation for children if there is problem of child's behavior and cooperation (World Health Organization (WHO), 2016). This show that actually the starting point of caries prevention is by making sure that mother or the parents are having knowledge of dental and oral hygiene to promote good dental and oral hygiene to their own children. Since parental knowledge has significant influence on children's dental caries (Isong, et al., 2012).

There is no discussion related with the finding...children with poor dental hygiene is more than good hygiene at parent with good parental knowledge ...why..??

The biggest percentage is poor parental knowledge and poor dental hygiene...why..???

Children with special healthcare needs (CSHCN) require health and related services more than that of children in general (McPherson et al., 1998). The children with special needs can suffer from mental and physical disabilities that influence them in mobility hindrance which resulting in higher poor dental hygiene (Bayarsaikhan, et al., 2015). This is in line with our study according to table 5 that either the parent have good knowledge, the highest category in dental and oral hygiene is poor. However, this is in accordance to the parental knowledge, which is in line with our study result in table 6 that there is relationship between parental knowledge and the dental and oral hygiene status of the children with special needs. Therefore, it is recommended that assessment of parental knowledge is important to

**Commented [a14]:** Should be 100% in each row.. Wrong in count the percentage...

**Commented [a15]:** Salah persentase nya...check dan hitung lagi

perform first before the appropriate education program for the target is held (Mahat & Bowen, 2017). The poor dental and oral hygiene of the children with special needs is also caused by poor routine dentist visit, perception that dental care is expensive (Hendaus et al., 2020), socioeconomic (Oberoi, et al., 2016), educational background (Al-batayneh, et al., 2019). The dental health personnel is needed to educate them, and it is recommended to initiate dental home process (Hendaus et al., 2020).

#### 4. Conclusion and Suggestion

We have presented a research to evaluate the relationship of parental knowledge and dental and oral hygiene status of the children with special needs in SDLB N 1 Martapura, Banjar Regency. The results comes into conclusion that there is a relationship between parental knowledge and dental and oral hygiene status of the children. It is shown by the results that the poor knowledge of the parent does in line with the poor dental and oral hygiene status of the children. We suggest that further research should be undertaken in the areas of factors related to the parental knowledge, so that the educational program held by dental personnel or Public Health Service is right on the target.

#### 5. Acknowledgments

We gratefully acknowledge the help provided by Poltekkes Kemenkes Banjarmasin, the Head of Education South Kalimantan Province, and SDLB N 1 Martapura, Banjar Regency for the cooperation and support.

#### 6. References

- Abdat, M., & Ramayana, I. (2020). Relationship between mother ' s knowledge and behaviour with oral health status of early childhood. *Padjadjaran Journal of Dentistry*, 32(3), 166-173. <https://doi.org/10.24198/pjd.vol32no2.24734>
- Al-batayneh, O. B., Al-khateeb, H. O., Ibrahim, W. M., & Khader, Y. S. (2019). Parental Knowledge and Acceptance of Different Treatment Options for Primary Teeth Provided by Dental Practitioners. *Frontiers in Public Health*, 7(November), 1-8. <https://doi.org/10.3389/fpubh.2019.00322>
- Alhabdan, Y. A., Albeshr, A. G., & Yenugadhathi, N. (2018). Prevalence of dental caries and associated factors among primary school children : a population-based cross-sectional study in Riyadh , Saudi Arabia. *Environmental Health and Preventive Medicine*, 23(60), 1-14.
- Bayarsaikhan, Z., Cruz, S., Neff, J., & Chi, D. L. (2015). Transitioning from pediatric care to adult care for adolescents with special health care needs: dentist perspectives (part 2). *Pediatr Dent.*, 37(5), 447-451.
- Bozorgmehr, E., Hajizamani, A., & Mohammadi, T. M. (2013). Oral Health Behavior of Parents as a Predictor of Oral Health Status of Their Children. *ISRN Dentistry*, (741783), 1-5. <https://doi.org/10.1155/2013/741783>
- Duijster, D., Jong-lenters, M. De, Verrips, E., & Loveren, C. Van. (2015). Establishing oral health promoting behaviours in children - parents ' views on barriers , facilitators and professional support : a qualitative study. *BMC Oral Health*, 15(157), 1-13. <https://doi.org/10.1186/s12903-015-0145-0>
- Greene, J. G., & Vermillion, J. R. (1964). The Simplified Oral Hygiene Index. *The Journal of The American Dental Association*, 68(1), 7-13. <https://doi.org/10.14219/jada.archive.1964.0034>
- Hendaus, M. A., Shaltout, D., Yasrab, D., Al-noubani, A., Hamad, S. G., Alamri, M., & Alhammadi, A. H. (2020). Parental Perception of a Dental Home for Children with Special Needs. *Pediatric Health, Medicine and Therapeutics*, 11, 379-384.
- Isong, I. A., Luff, D., Perrin, J. M., Winickoff, J. P., & Ng, M. W. (2012). Parental Perspectives of Early Childhood Caries. *Clinical Pediatrics*, 51(1), 77-85. <https://doi.org/10.1177/0009922811417856>
- Jürgensen, N., & Petersen, P. E. (2009). Oral health and the impact of socio-behavioural factors in a cross sectional survey of 12-year old school children in Laos. *BMC Oral Health*, 9(29), 1-11. <https://doi.org/10.1186/1472-6831-9-29>
- Kubota, Y., Jareonpituak, S., Narksawat, K., Satitvipawee, P., & Durward, C. (2017). Status , knowledge , attitudes , and practices of 12-18-year old orphaned children pertaining to dental caries in an dental clinic by non-governmental organization at Phnom. *Oral Health Care*, 2(2), 1-4. <https://doi.org/10.15761/OHC.1000117>

Commented [m16]: Check all OUT OF DATE reference

- Lai, B., Tan, W. K., & Lu, Q. S. (2018). Clinical efficacy of a two-year oral health programme for infants and toddlers in Singapore. *Singapore Med J*, 59(2), 87-93. <https://doi.org/10.11622/smedj.2017045>
- Maharani, D. A., Zhang, S., Gao, S. S., & Chu, C. (2019). Dental Caries and the Erosive Tooth Wear Status of 12-Year-Old Children in Jakarta, Indonesia. *International Journal of Environmental Research and Public Health*, 16(2994), 1-10. <https://doi.org/10.3390/ijerph16162994>
- Mahat, G., & Bowen, F. (2017). Parental Knowledge about Urban Preschool Children's Oral Health Risk. *Pediatric Nursing*, 43(1), 30-34.
- Mchugh, M. L. (2013). The Chi-square test of independence. *Biochemia Medica*, 23(2), 143-149. <https://doi.org/10.11613/BM.2013.018>
- McPherson, M., Arango, P., Fox, H., Lauver, C., McManus, M., Newacheck, P. W., ... Strickland, B. (1998). A New Definition of Children With Special Health Care Needs. *Pediatrics*, 102(1), 137-140.
- Ministry of Health of Indonesia. (2019). Faktor Risiko Kesehatan Gigi dan Mulut. *InfoDATIN*, 1-6. Retrieved from <https://pusdatin.kemkes.go.id/download.php?file=download/pusdatin/infodatin/infodatin-gigi.pdf>
- Mota-veloso, I., Nogueira, R., Eduardo, C., Alca, P., Aparecida, S., Ramos-jorge, M. L., & Oliveira-ferreira, F. (2012). Research in Developmental Disabilities Does intellectual disability affect the development of dental caries in patients with cerebral palsy? *Research in Developmental Disabilities*, 33, 1503-1507. <https://doi.org/10.1016/j.ridd.2012.03.026>
- Ningrum, V., Bakar, A., Shieh, T., & Shih, Y. (2021). The Oral Health Inequities between Special Needs Children and Normal Children in Asia : A Systematic Review and. *Healthcare*, 9(410), 1-12. <https://doi.org/10.3390/healthcare9040410>
- Oberoi, S. S., Sharma, G., & Oberoi, A. (2016). A cross-sectional survey to assess the effect of socioeconomic status on the oral hygiene habits. *J Indian Soc Periodontol*, 20(5), 531-542. <https://doi.org/10.4103/0972-124X.201629>
- Pini, D. D. M., Fröhlich, P. C. G. R., & Rigo, L. (2016). Oral health evaluation in special needs individuals. *Einstein (Sao Paulo)*, 14(55 54), 501-507. <https://doi.org/10.1590/S1679-45082016AO3712>
- Priya, M. M., Kumar, P., & Saraswathi, S. (2018). Assessment of parental oral health knowledge, attitude, and behavior. *Journal of Pharmacy Research*, 12(1), 10-13.
- Rogéria, A., Castilho, F. De, Luiz, F., Souza, T. De, & Puppim-rontani, R. M. (2013). Influence of family environment on children's oral health : a systematic review ☆. *Journal de Pediatria*, 89(2), 116-123. <https://doi.org/10.1016/j.jpmed.2013.03.014>
- Sarvas, E. (2017). Assessing Caries Risk in Children With Special Needs. *Decisions in Dentistry*, 3(3), 48-52.
- Syreen, S., Anwar, A., Ahmad, G., & Rahman, M. (2018). Prevalence of Dental Caries among Children Residing in Muslim Dominated. *Journal of Medical Science and Clinical Research*, 06(12), 817-819.
- Urwannachotima, N., & Hanvoravongchai, P. (2020). System dynamics analysis of dental caries status among Thai adults and elderly. *Journal of Health Research*, 34(2), 134-146. <https://doi.org/10.1108/JHR-05-2019-0097>
- Vellappally, S., Gardens, S. J., Kheraif, A. A. Al, Krishna, M., Babu, S., Hashem, M., ... Anil, S. (2014). The prevalence of malocclusion and its association with dental caries among 12-18-year-old disabled adolescents. *BMC Oral Health*, 14(123), 1-7.
- World Health Organization. (2003). *Continuous improvement of oral health in the 21st century - the approach of the WHO Global Oral Health Programme*. Geneva.
- World Health Organization (WHO). (2016). *WHO Expert Consultation on Public Health Intervention against Early Childhood Caries*. Retrieved from <https://apps.who.int/iris/bitstream/handle/10665/255627/WHO-NMH-PND-17.1-eng.pdf>
- Yeung, P. M., Wong, V. C. N., Mcgrath, C. P., Yiu, C. K. Y., & Lee, G. H. M. (2019). Oral health status of children with epilepsy in Hong Kong. *J Invest Clin Dent*, 10(February), 1-8. <https://doi.org/10.1111/jicd.12479>
- Zhang, S., Chau, A. M. H., Lo, E. C. M., & Chu, C. (2014). Dental caries and erosion status of 12-year-old Hong Kong children. *BMC Public Health*, 14(7), 1-7.

Jurnal Riset Kesehatan, x (x), xxxx, 7 - 7

DOI: 10.31983/jrk.vxix.xxxx



---

## PARENTAL KNOWLEDGE TOWARDS CHILDREN'S DENTAL AND ORAL HYGIENE IN SDLB 1 MARTAPURA

**Agno Elman Efzi ; Emilda Sari\***

*Department of Dental Hygiene, Poltekkes Kemenkes Banjarmasin ; Jl. Mistar Cokrokusumo No.1A ; Banjarbaru 70714 ; Indonesia*

### Abstract

Dental and oral hygiene problem is the leading cause of daily activities disruption in children such as missing school, bad learning concentration, appetite and nutritional intake. Knowledge of dental and oral hygiene should be given at early age in family by the parents. The aim of our work was to determine relationship between parents' knowledge and dental hygiene of the SDLB children (children with special needs). We undertook this research with analytical survey. A cross-sectional design was carried out. The sampling method was total sampling namely parents of children with special needs and the children as many as 80 people. Chi-Square test was performed to analyze the data. The results show that the knowledge of parents about dental and oral hygiene is not good (85%), while the dental and oral hygiene in children with special needs is poor (85%). Based on the Chi-Square Test, it produces a p-value of sig (2-sided) = 0.002 in the test, so it can obtain a p-value (probability value) from the test ( $p = 0.002 < = 0.05$ ) so that  $H_0$  is rejected. In conclusion, a relationship between parental knowledge and oral hygiene in children with special needs does exist.

**Keywords:** *knowledge; dental and oral hygiene; children with special needs; parents*

### 1. Introduction

Caries is the most common chronic dental problem yet it is actually preventable (Syreen, et al. 2018; Alhabdan, et al., 2018). Caries attacked 3.5 billion people in the world and it is started when the first teeth appear at children age. The report of World Health Organization reveals 60 - 90% of school children worldwide has experienced caries especially in Asian

countries (Van Chuyen, et al., 2021). A study found that the caries prevalence and DMFT score of a group of children aged 12 years old in Indonesia are higher than children in the same age in Southeast Asian countries (Zhang, et al. 2014; Urwannachotima & Hanvoravongchai, 2020; Lai, et al., 2018; Kubota, et al., 2017). It is proven that 61% of 12-year-old Indonesian children were affected by dental caries with the mean DMFT score in 1.58 (Maharani, et al., 2019).

Children may experience psychiatric and mental health issues. Special needs children are those under the age of 18 who have either

---

\*) Corresponding Author (Emilda Sari)

E-mail: [melda\\_akg\\_bjm@yahoo.co.id](mailto:melda_akg_bjm@yahoo.co.id)

physical or cognitive disabilities including intellectual disability (ID), Down Syndrome (DS), autism spectrum disorder (ASD), and attention deficit hyperactivity disorder (ADHD). They have higher risk of dental problems and poor oral hygiene than normal children (Yeung, et al., 2019; Mota-veloso et al., 2012; Vellappally et al., 2014). This condition can lead to dental aversion which associated with increase plaque levels and caries experience. Intellectual disabilities may also limit the self - care, resulting in poor dental hygiene (Sarvas, 2017). Children with special needs require more assistance because of their mental as well as physical challenges, even if they are over seven years old. Because some of them may learn slowly and being uncooperative, have difficulty to understand the behavior of dental hygiene such as brushing teeth (Ningrum, et al., 2021). It was mentioned that children with special need is prone having a high dental carries index and inadequate oral hygiene (Pini, et al., 2016).

The role of parents is essential in dental hygiene as they are the main caregivers to their children. The behavior of parents, particularly mothers can affect their children’s health. Some of the behavior of the parents include tooth brushing habit, dietary, and food choices. A good behavior of the parents can affect the children in the effort of caries risk reduction(Bozorgmehr, et al., 2013). Family environment has something to do with the influence of good behavior about dental hygiene(Duijster, et al., 2015).

The relationship of the parental knowledge regarding the dental hygiene of their children are very important to understand. Over the past ten years, the information of the relationship of the parent on the dental hygiene of the children have been conducted. A significant relationship of parental behavior and preschool children in Kerman, South East of Iran was proven in

(Bozorgmehr et al., 2013). However, this study doesn’t evaluate the knowledge of the parent among the relationship to the children dental hygiene, though it was mentioned that knowledge and attitude of the parents affect children dental health behavior and status. A survey on parental oral knowledge and behavior has been conducted in relationship to the children oral health in (Priya, et al., 2018; Abdat & Ramayana, 2020; Rogéria, et al., 2013). The results indicate that the children poor oral health practice is reflected by the parent partial knowledge and behavior toward oral health. However, all aforementioned research suffer from a relationship to the children with special needs.

Our objective was to evaluate the relationship of parental knowledge to the dental hygiene of the children with special needs. The study was performed by evaluating the knowledge of the parents by using questionnaire and the dental oral hygiene by examining OHI-S of the children. A further objective was analyzing the parents’ knowledge and the dental hygiene of the children with special needs.

## 2. Method

### Study design and sampling procedure

Our study type was analytical survey. The cross-sectional approach was used to determine the relationship between parental knowledge and oral hygiene and the approach and the variables of research were obtained in the same time. The study was conducted from October 2019 to May 2020. The variable of research consists of independent variable, i.e. knowledge of parents and dependent variable i.e. dental and oral hygiene of the children. Table 1 shows the questionnaire of knowledge for the parents regarding dental and oral health.

**Table 1.** Questionnaire of knowledge

| Question   | Multiple choice  |
|--|--|
| How many times a day at least brush your teeth?              | a. One time<br>b. Two times<br>c. Four times   |
| When is the right time to brush your teeth?                  | a. Morning shower<br>b. Morning and evening shower<br>c. After breakfast and before going to bed |
| How many times should we check our teeth and mouth at least? | a. Once a year<br>b. Once in six months  |



|   |   |
|---|---|
|   | c. Tooth brushing is enough   |
| How do you brush the back/chewing part of your teeth?                 | a. Back and forth<br>b. Prying<br>c. Up and down  |
| What are the signs that there is a problem with oral health?          | a. Pure white teeth<br>b. No bad breath<br>c. Gums bleed when brushing  |
| Which food is not good for oral health?                               | a. Sweet and sticky food<br>b. Fibrous and watery foods<br>c. Foods that contain vitamin C  |
| One of the consequences of not taking care of dental and oral health? | a. Cavities<br>b. Fresh breath<br>c. Strong teeth   |
| How to brush your teeth properly and correctly?                       | a. Slowly and gently<br>b. Quickly<br>c. Quickly and hard   |
| Is it okay to share a toothbrush with other people?                   | a. Yes, if you know each other<br>b. Yes, if you are family<br>c. Should not  |
| How to chew food properly?  | a. Both sides of jaw<br>b. Left jaw only<br>c. Right jaw only   |
| What is the purpose of tooth brushing?                                | a. Cleaning teeth from food residue and massage the gum<br>b. Massage the gum<br>c. Whitening the teeth   |
| What should you use when brushing your teeth?                         | a. Siwak<br>b. Mouth wash<br>c. Tooth paste   |
| What are the requirements for a good toothbrush?                      | a. A toothbrush with a slender brush head and soft bristles<br>b. Curved toothbrush and uneven bristles<br>c. Toothbrush with straight handle and coarse bristles |
| What are the foods that clean teeth?                                  | a. Fibrous food<br>b. Soft food<br>c. Hard food   |
| What is the purpose of dental and oral hygiene?                       | a. Prevents infections of the mouth, teeth and tongue<br>b. Makes bad breath<br>c. Gives you a feeling of insecurity  |
| Which of the following factors are said to be healthy oral?           | a. Healthy dental support tissue<br>b. There is tartar in the teeth<br>c. Bleeding gums when brushing teeth   |
| What a tooth paste should contain to be said as a good tooth paste?   | a. Foam<br>b. Fluor<br>c. Perfume   |
| What problem caused by tooth decay?                                   | a. Problems with growth and development of children   |

|                                       |   |
|---------------------------------------|---|
|                                       | b. Make children more appetizing<br>c. Keeping children's dental and oral health                    |
| What is the effect of dental problem? | a. A good nutrition absorption<br>b. Decrease appetite<br>c. Perfect breakdown of food in the mouth |

The population was all 40 students with special needs and 40 parents at SDLB N 1 Martapura, Banjar Regency. Total sampling was carried out with the total population, namely all students with special needs and their parents at SDLB 1 Martapura, Banjar Regency with a total sample of 80 people.

**Data collection**

Primary data collection was carried out by conducting direct examinations of children with

special needs at the SDLB N 1 Martapura, Banjar Regency using a diagnostic instrument set, OHI-S format and giving a dental and oral hygiene knowledge questionnaire to parents of children with special needs. The questionnaire question was 19 in total as shown in Table 1. The validity of question has been done using Corrected-Item Total-Collection. The results of Corrected-Item Total-Collection is shown in Table 2.

**Table 2.** Item-Total Statistics

|     | Scale Mean if Item Deleted | Scale Variance if Item Deleted | Corrected Item-Total Correlation | Cronbach's Alpha If Item Deleted |
|-----|----------------------------|--------------------------------|----------------------------------|----------------------------------|
| p1  | 11.73                      | 62.202                         | .843                             | .968                             |
| p2  | 11.67                      | 63.954                         | .622                             | .971                             |
| p3  | 11.63                      | 62.516                         | .819                             | .969                             |
| p6  | 11.63                      | 63.895                         | .637                             | .971                             |
| p7  | 11.67                      | 62.575                         | .801                             | .969                             |
| p8  | 11.60                      | 62.869                         | .786                             | .969                             |
| p9  | 11.73                      | 62.202                         | .843                             | .968                             |
| p11 | 11.67                      | 62.575                         | .801                             | .969                             |
| p12 | 11.73                      | 62.202                         | .843                             | .968                             |
| p13 | 11.60                      | 63.490                         | .703                             | .970                             |
| p15 | 11.60                      | 62.869                         | .786                             | .969                             |
| p16 | 11.67                      | 62.575                         | .801                             | .969                             |
| p17 | 11.67                      | 62.575                         | .801                             | .969                             |
| p18 | 11.60                      | 63.490                         | .703                             | .970                             |
| p19 | 11.63                      | 62.516                         | .819                             | .969                             |
| p20 | 11.60                      | 62.869                         | .786                             | .969                             |
| p21 | 11.73                      | 62.202                         | .843                             | .968                             |
| p22 | 11.67                      | 62.575                         | .801                             | .969                             |
| p23 | 11.60                      | 62.869                         | .786                             | .969                             |
| p24 | 11.63                      | 63.895                         | .637                             | .971                             |
| p25 | 11.60                      | 63.490                         | .703                             | .970                             |

According to table 2, the correlation is taken from Corrected Item-Total Correlation. The values are compared to r table. Because the total

item is 21, so the r table value is  $21-2 = 19$ . The r table is in significance 0.05 or 5%. The value is 0.4329(Kamilah, 2015). Because all the values in Corrected Item-Total Correlation is more than

0.4329, so, it can be concluded that the questions are valid.

Each of question consists of 3 answers. Score 1 is given to the correct answer and score 0 is given to the incorrect answer. There are two categories in knowledge, i.e. Good if the score is 10 – 19, and poor if the score is 0 – 9.

Secondary data was data obtained from the research site, SDLB N 1 Martapura, Banjar Regency, regarding the names of students and their parents, age, gender, type of disability and the number of students with special needs and the parents.

### Data analysis

Data analysis was started by inputting the primary and secondary in frequency distribution table. After collecting the results of dental and oral hygiene examinations for children with special needs, then the results were statistically tested using the SPSS program with the Chi Square test to find out the existence of relationship in parental knowledge and oral hygiene (Mchugh, 2013). The category used for Debris Index and Calculus is based on in which good is scored 0 – 0.6; fair if scored 0.7 – 1.8; poor if 1.9 – 3.0 (Greene & Vermillion, 1964).

### 3. Result and Discussion

Results on frequency distribution are divided into distribution based on the parents and children. The data of parents include the number of respondents and knowledge. Meanwhile the data of children includes the number of respondents, and dental and oral hygiene.

**Table 3.** Frequency distribution of parent respondents

| Gender | Number of respondents | Percentage (%) |
|--------|-----------------------|----------------|
| Male   | 12                    | 30             |
| Female | 28                    | 70             |
| Total  | 40                    | 100            |

Table 3 shows the number of parents of children with special needs in which 30% of the respondents are male parent (father) and 70% are female parent (mother). In this research, not all of

the respondents are mothers, because not all of the children go to school with their mothers. Some of them are accompanied by their fathers. Even their fathers wait and pick up the children. Thus, the researchers found difficulty to meet the mother of each children. Some of the parents are also comfortable if the respondent is the father.

**Table 4.** Frequency distribution of students in SDLB N 1 Martapura

| Grade     | Number | Percentage (%) |
|-----------|--------|----------------|
| Grade I   | 12     | 30             |
| Grade II  | 10     | 25             |
| Grade III | 5      | 12.5           |
| Grade IV  | 4      | 10             |
| Grade V   | 4      | 10             |
| Grade VI  | 5      | 12.5           |
| Total     | 40     | 100            |

Table 4 shows the frequency distribution of the students that the total number of students in SDLB N 1 Martapura is 40.

**Table 5.** Frequency distribution of parental knowledge

| Category | Number of respondents | Percentage (%) |
|----------|-----------------------|----------------|
| Good     | 6                     | 15             |
| Poor     | 34                    | 85             |
| Total    | 40                    | 100            |

Table 5 states the number of parental knowledge with poor category is 34 people and good category is 6 people.

**Table 6.** Frequency distribution of dental and oral hygiene of the children

| Category | Number of Respondents | Percentage (%) |
|----------|-----------------------|----------------|
| Good     | 4                     | 10             |
| Fair     | 2                     | 5              |
| Poor     | 34                    | 85             |
| Total    | 40                    | 100            |

Table 6 shows the highest dental and oral hygiene percentage is in category of poor with 34 children or 85% of the total respondents.

**Table 7.** Cross Tabulation of the Relationship between Parental Knowledge and Dental and Oral Hygiene in Children with Special Needs

| Parental know-ledge | Dental and oral hygiene of the children |      |      |      |      |      | Total |     |
|---------------------|---|------|------|------|------|------|-------|-----|
|                     | Good                                    |      | Fair |      | Poor |      | n     | %   |
|                     | n                                       | %    | n    | %    | n    | %    |       |     |
| Good                | 1                                       | 16.7 | 2    | 33.3 | 3    | 50   | 6     | 100 |
| Poor                | 3                                       | 8.8  | 0    | 0    | 31   | 91.2 | 34    | 100 |
| Total               | 4                                       | 10   | 2    | 5    | 34   | 85   | 40    | 100 |

Table 7 shows from 40 parents, at good parental knowledge, there is 1 parent (16.7%) who has children with good dental and oral hygiene, 2 parents (33.3%) who have children with fair dental and oral hygiene and 3 parents (50%) who have children with poor dental and oral hygiene.

Meanwhile, at poor parental knowledge, there are 3 parents (8.8%) who have children with good dental and oral hygiene and 31 parents (91.2%) who have children with poor dental and oral hygiene.

**Table 8.** Chi Square test results on parental knowledge and dental oral hygiene of the children

|                   | Value               | Df | Sig (2-sided) |
|-------------------|---------------------|----|---------------|
| Person Chi-Square | 12.664 <sup>a</sup> | 2  | 0.002         |

Table 8 shows the p value in the sig (2-sided) = 0.002 column in the test, meaning that the p value (probability value) of the test is less than, so  $H_0$  is rejected. In conclusion, there is a relationship between parental knowledge and dental and oral hygiene in children with special needs at SDLB N 1 Martapura. So, if the parental knowledge about dental and oral health is good, the dental and oral hygiene of their children can be way better too.

In table 7, the number of parents who have poor knowledge of dental and oral hygiene is higher than those who have good poor knowledge. As shown in the research of (Mahat & Bowen, 2017), parents actually have awareness that teeth are important to maintain. However few understand the cavities appear in their children's teeth which cause cavities in

permanent teeth. So, the bad dental and oral hygiene may be caused by baby teeth cavities that parents didn't realize.

One objective of Ministry of Health of Indonesia 2030 is that to reduce the proportion of dental caries experience of children at age 12 (Ministry of Health of Indonesia, 2019). WHO recommends three prevention of childhood caries i.e. primary prevention which includes primary health care programs especially for maternal and child health that is conducted in early age such as no sugars for baby until 2 years. Secondary prevention is performed focusing on early detection of carious lesions. This is focusing on the dental personnel and even mothers to detect early signs of carious lesions. Tertiary prevention includes reduce cavity by avoiding unnecessary extraction and restoration function. In tertiary prevention may apply rehabilitation for children if there is problem of child's behavior and cooperation (World Health Organization (WHO), 2016). This show that actually the starting point of caries prevention is by making sure that mother or the parents are having knowledge of dental and oral hygiene to promote good dental and oral hygiene to their own children. Since parental knowledge has significant influence on children's dental caries (Isong, et al., 2012).

In table 7, it shows that at parent with good parental knowledge, the number of children who have poor dental and oral hygiene is higher than that with good dental and oral hygiene. This shows that even though parents have good knowledge, however, most children with special needs have less awareness in keeping dental and oral health so that there is still a lot of those who have poor dental and oral hygiene. Children with special needs are at higher risk that normal children because they may have impaired cognitive abilities, behavioral problems, poor motor coordination, uncontrolled body movements, neuromuscular problem such as drooling, swallowing problem (Ningrum et al., 2021). These problems can hamper them to clean their own teeth or use the common brushing method. Besides, the problems can also reduce the saliva flow that naturally can help wash the food residue in their mouth (Zamani, 2010).

Table 7 shows that the biggest percentage is poor parental knowledge and poor dental hygiene of the children in 91.2%. The biggest obstacle faced by most of parents regarding dental services to their children is the awareness. Parents tend to visit dentist when there is necessity and not experience professional dental care. This can be caused by the lack of information given to parents regarding oral health from health and social care staffs. Besides, the level of education can also be the factor of poor knowledge of the parents about dental and oral care to their children. It is mentioned that stigma of the shame having a child with special needs also becomes a factor that parents having fear of being discriminated (Hegde et al., 2015). Thus, parents can be isolated, not getting any support and information which enable them to access oral health care. In consequence, their knowledge is poor. This situation impact to the children that the children with special needs don't get any dental care service at home.

Children with special healthcare needs (CSHCN) require health and related services more than that of children in general. The children with special needs can suffer from mental and physical disabilities that influence them in mobility hindrance which resulting in higher poor dental hygiene (Bayarsaikhan, et al., 2015). This is in line with our study according to table 7 that even the parent have good knowledge, the highest category in dental and oral hygiene is poor. However, this is in accordance to the parental knowledge, which is in line with our study result in table 8 that there is relationship between parental knowledge and the dental and oral hygiene status of the children with special needs. Therefore, it is recommended that assessment of parental knowledge is important to perform first before the appropriate education program for the target is held (Mahat & Bowen, 2017). The poor dental and oral hygiene of the children with special needs is also caused by poor routine dentist visit, perception that dental care is expensive (Hendaus et al., 2020), socioeconomic (Oberoi, et al., 2016), educational background (Albatayneh, et al., 2019). The dental health personnel is needed to educate them, and it is recommended to initiate dental home process (Hendaus et al., 2020).

#### 4. Conclusion and Suggestion

We have presented a research to evaluate the relationship of parental knowledge and dental and oral hygiene status of the children with special needs in SDLB N 1 Martapura, Banjar Regency. The results comes into conclusion that there is a relationship between parental knowledge and dental and oral hygiene status of the children. It is shown by the results that the poor knowledge of the parent does in line with the poor dental and oral hygiene status of the children. Even though parents have good knowledge, however, most children with special needs have less awareness in keeping dental and oral health so that there is still a lot of those who have poor dental and oral hygiene. It is due to the barriers such as behavioral problem experienced by children with special needs. We suggest that the parents engage more in the importance of dental and oral hygiene and follow the instruction from health practitioners regarding their children dental and oral health. Besides, Public Health Center can increase the education of dental and oral health towards the parents in SDLB so that parental knowledge can be better.

#### 5. Acknowledgments

We gratefully acknowledge the help provided by Poltekkes Kemenkes Banjarmasin, the Head of Education South Kalimantan Province, and SDLB N 1 Martapura, Banjar Regency for the cooperation and support.

#### 6. References

- Abdat, M., & Ramayana, I. (2020). Relationship between mother ' s knowledge and behaviour with oral health status of early childhood. *Padjadjaran Journal of Dentistry*, 32(3), 166-173. <https://doi.org/10.24198/pjd.vol32no2.24734>
- Al-batayneh, O. B., Al-khateeb, H. O., Ibrahim, W. M., & Khader, Y. S. (2019). Parental Knowledge and Acceptance of Different Treatment Options for Primary Teeth Provided by Dental Practitioners. *Frontiers in Public Health*, 7(November), 1-8. <https://doi.org/10.3389/fpubh.2019.00322>
- Alhabdan, Y. A., Albeshr, A. G., & Yenugadhathi, N. (2018). Prevalence of dental caries and associated factors among primary school

- children : a population-based cross-sectional study in Riyadh , Saudi Arabia. *Environmental Health and Preventive Medicine*, 23(60), 1-14.
- Bayarsaikhan, Z., Cruz, S., Neff, J., & Chi, D. L. (2015). Transitioning from pediatric care to adult care for adolescents with special health care needs: dentist perspectives (part 2). *Pediatr Dent.*, 37(5), 447-451.
- Bozorgmehr, E., Hajizamani, A., & Mohammadi, T. M. (2013). Oral Health Behavior of Parents as a Predictor of Oral Health Status of Their Children. *ISRN Dentistry*, (741783), 1-5. <https://doi.org/10.1155/2013/741783>
- Duijster, D., Jong-lenters, M. De, Verrips, E., & Loveren, C. Van. (2015). Establishing oral health promoting behaviours in children – parents’ views on barriers , facilitators and professional support : a qualitative study. *BMC Oral Health*, 15(157), 1-13. <https://doi.org/10.1186/s12903-015-0145-0>
- Greene, J. G., & Vermillion, J. R. (1964). The Simplified Oral Hygiene Index. *The Journal of The American Dental Association*, 68(1), 7-13. <https://doi.org/10.14219/jada.archive.1964.0034>
- Hegde, A. M., Babu, A. A., Mohammed, A., John, A., Singh, K., C., P. V., & Shetty, S. (2015). Special Needs of Special Children-Parental View. *Journal of Health and Allied Sciences NU*, 05(02), 038-044. <https://doi.org/10.1055/s-0040-1703887>
- Hendaus, M. A., Shaltout, D., Yasrab, D., Al-noubani, A., Hamad, S. G., Alamri, M., & Alhammadi, A. H. (2020). Parental Perception of a Dental Home for Children with Special Needs. *Pediatric Health, Medicine and Therapeutics*, 11, 379-384.
- Isong, I. A., Luff, D., Perrin, J. M., Winickoff, J. P., & Ng, M. W. (2012). Parental Perspectives of Early Childhood Caries. *Clinical Pediatrics*, 51(1), 77-85. <https://doi.org/10.1177/0009922811417856>
- Kamilah, E. N. (2015). *DISTRIBUSI NILAI r. Pengaruh Keterampilan Mengajar Guru Terhadap Hasil Belajar Siswa Pada Mata Pelajaran Akuntansi Universitas*, 91.
- Kubota, Y., Jareonpituak, S., Narksawat, K., Satitvipawee, P., & Durward, C. (2017). Status , knowledge , attitudes , and practices of 12-18-year old orphaned children pertaining to dental caries in an dental clinic by non-governmental organization at Phnom. *Oral Health Care*, 2(2), 1-4. <https://doi.org/10.15761/OHC.1000117>
- Lai, B., Tan, W. K., & Lu, Q. S. (2018). Clinical efficacy of a two-year oral health programme for infants and toddlers in Singapore. *Singapore Med J*, 59(2), 87-93. <https://doi.org/10.11622/smedj.2017045>
- Maharani, D. A., Zhang, S., Gao, S. S., & Chu, C. (2019). Dental Caries and the Erosive Tooth Wear Status of 12-Year-Old Children in Jakarta , Indonesia. *International Journal of Environmental Research and Public Health*, 16(2994), 1-10. <https://doi.org/10.3390/ijerph16162994>
- Mahat, G., & Bowen, F. (2017). Parental Knowledge about Urban Preschool Children’s Oral Health Risk. *Pediatric Nursing*, 43(1), 30-34.
- Mchugh, M. L. (2013). The Chi-square test of independence. *Biochemia Medica*, 23(2), 143-149. <https://doi.org/10.11613/BM.2013.018>
- Ministry of Health of Indonesia. (2019). Faktor Risiko Kesehatan Gigi dan Mulut. *InfoDATIN*, 1-6. Retrieved from <https://pusdatin.kemkes.go.id/download.php?file=download/pusdatin/infodatin/infodatin-gigi.pdf>
- Mota-veloso, I., Nogueira, R., Eduardo, C., Alca, P., Aparecida, S., Ramos-jorge, M. L., & Oliveira-ferreira, F. (2012). Research in Developmental Disabilities Does intellectual disability affect the development of dental caries in patients with cerebral palsy ? *Research in Developmental Disabilities*, 33, 1503-1507. <https://doi.org/10.1016/j.ridd.2012.03.026>
- Ningrum, V., Bakar, A., Shieh, T., & Shih, Y. (2021). The Oral Health Inequities between Special Needs Children and Normal Children in Asia : A Systematic Review and. *Healthcare*, 9(410), 1-12. <https://doi.org/10.3390/healthcare9040410>
- Oberoi, S. S., Sharma, G., & Oberoi, A. (2016). A cross-sectional survey to assess the effect of socioeconomic status on the oral hygiene habits. *J Indian Soc Periodontol*, 20(5), 531-542. <https://doi.org/10.4103/0972-124X.201629>
- Pini, D. D. M., Fröhlich, P. C. G. R., & Rigo, L. (2016). Oral health evaluation in special needs individuals. *Einstein (Sao Paulo)*, 14(55 54), 501-507. <https://doi.org/10.1590/S1679->

- 45082016AO3712
- Priya, M. M., Kumar, P., & Saraswathi, S. (2018). Assessment of parental oral health knowledge , attitude , and behavior. *Journal of Pharmacy Research*, 12(1), 10–13.
- Rogéria, A., Castilho, F. De, Luiz, F., Souza, T. De, & Puppini, R. M. (2013). Influence of family environment on children ' s oral health : a systematic review ☆. *Journal de Pediatria*, 89(2), 116–123. <https://doi.org/10.1016/j.jpmed.2013.03.014>
- Sarvas, E. (2017). Assessing Caries Risk in Children With Special Needs. *Decisions in Dentistry*, 3(3), 48–52.
- Syreen, S., Anwar, A., Ahmad, G., & Rahman, M. (2018). Prevalence of Dental Caries among Children Residing in Muslim Dominated. *Journal of Medical Science and Clinical Research*, 06(12), 817–819.
- Urwannachotima, N., & Hanvoravongchai, P. (2020). System dynamics analysis of dental caries status among Thai adults and elderly. *Journal of Health Research*, 34(2), 134–146. <https://doi.org/10.1108/JHR-05-2019-0097>
- Van Chuyen, N., Van Du, V., Van Ba, N., Long, D. D., & Son, H. A. (2021). The prevalence of dental caries and associated factors among secondary school children in rural highland Vietnam. *BMC Oral Health*, 21(1), 1–7. <https://doi.org/10.1186/s12903-021-01704-y>
- Vellappally, S., Gardens, S. J., Kheraif, A. A. Al, Krishna, M., Babu, S., Hashem, M., ... Anil, S. (2014). The prevalence of malocclusion and its association with dental caries among 12-18-year-old disabled adolescents. *BMC Oral Health*, 14(123), 1–7.
- World Health Organization (WHO). (2016). *WHO Expert Consultation on Public Health Intervention against Early Childhood Caries*. Retrieved from <https://apps.who.int/iris/bitstream/handle/10665/255627/WHO-NMH-PND-17.1-eng.pdf>
- Yeung, P. M., Wong, V. C. N., Mcgrath, C. P., Yiu, C. K. Y., & Lee, G. H. M. (2019). Oral health status of children with epilepsy in Hong Kong. *J Invest Clin Dent*, 10(February), 1–8. <https://doi.org/10.1111/jicd.12479>
- Zamani, A. R. (2010). Oral Health for Children with Disabilities and Special Needs. *California Childcare Health Program*, 1–2.
- Zhang, S., Chau, A. M. H., Lo, E. C. M., & Chu, C. (2014). Dental caries and erosion status of 12-year-old Hong Kong children. *BMC Public Health*, 14(7), 1–7.