

ISSN-0976-0245 (Print) • ISSN-0976-5506 (Electronic)

Volume 10

Number 1

January 2019



Indian Journal of Public Health Research & Development

An International Journal

SCOPUS IJPHRD CITATION SCORE

Indian Journal of Public Health Research and Development
Scopus coverage years: from 2010 to 2017 Publisher:
R.K. Sharma, Institute of Medico-Legal Publications
ISSN:0976-0245E-ISSN: 0976-5506 Subject area: Medicine:
Public Health, Environmental and Occupational Health
CiteScore 2015- 0.02
SJR 2015- 0.105
SNIP 2015- 0.034



Website:

www.ijphrd.com

Indian Journal of Public Health Research & Development

EXECUTIVE EDITOR

Prof Vidya Surwade

Associate Professor, Dr Baba Saheb Ambedkar, Medical College & Hospital, Rohinee, Delhi

INTERNATIONAL EDITORIAL ADVISORY BOARD

1. **Dr. Abdul Rashid Khan** B. Md Jagar Din, (*Associate Professor*)
Department of Public Health Medicine, Penang Medical College, Penang, Malaysia
2. **Dr. V Kumar** (*Consulting Physician*)
Mount View Hospital, Las Vegas, USA
3. **Basheer A. Al-Sum**,
Botany and Microbiology Deptt, College of Science, King Saud University,
Riyadh, Saudi Arabia
4. **Dr. Ch Vijay Kumar** (*Associate Professor*)
Public Health and Community Medicine, University of Buraimi, Oman
5. **Dr. VMC Ramaswamy** (*Senior Lecturer*)
Department of Pathology, International Medical University, Bukit Jalil, Kuala Lumpur
6. **Kartavya J. Vyas** (*Clinical Researcher*)
Department of Deployment Health Research,
Naval Health Research Center, San Diego, CA (USA)
7. **Prof. PK Pokharel** (*Community Medicine*)
BP Koirala Institute of Health Sciences, Nepal

NATIONAL SCIENTIFIC COMMITTEE

1. **Dr. Anju Ade** (*Associate Professor*)
Navodaya Medical College, Raichur, Karnataka
2. **Dr. E. Venkata Rao** (*Associate Professor*) Community Medicine,
Institute of Medical Sciences & SUM Hospital, Bhubaneswar, Orissa.
3. **Dr. Amit K. Singh** (*Associate Professor*) Community Medicine,
VCSG Govt. Medical College, Srinagar – Garhwal, Uttarakhand
4. **Dr. R G Viveki** (*Professor & Head*) Community Medicine,
Belgaum Institute of Medical Sciences, Belgaum, Karnataka
5. **Dr. Santosh Kumar Mulage** (*Assistant Professor*)
Anatomy, Raichur Institute of Medical Sciences Raichur (RIMS), Karnataka
6. **Dr. Gouri Ku. Padhy** (*Associate Professor*) Community and Family
Medicine, All India Institute of Medical Sciences, Raipur
7. **Dr. Ritu Goyal** (*Associate Professor*)
Anaesthesia, Sarswathi Institute of Medical Sciences, Panchsheel Nagar
8. **Dr. Anand Kalaskar** (*Associate Professor*)
Microbiology, Prathima Institute of Medical Sciences, AP
9. **Dr. Md. Amirul Hassan** (*Associate Professor*)
Community Medicine, Government Medical College, Ambedkar Nagar, UP
10. **Dr. N. Girish** (*Associate Professor*) Microbiology, VIMS&RC, Bangalore
11. **Dr. BR Hungund** (*Associate Professor*) Pathology, JNMC, Belgaum.
12. **Dr. Sartaj Ahmad** (Assistant Professor),
Medical Sociology, Department of Community Medicine, Swami Vivekananda Subharti
University, Meerut, Uttar Pradesh, India
13. **Dr Sumeeta Soni** (Associate Professor)
Microbiology Department, B.J. Medical College, Ahmedabad, Gujarat, India

NATIONAL EDITORIAL ADVISORY BOARD

1. **Prof. Sushanta Kumar Mishra** (Community Medicine)
GSL Medical College – Rajahmundry, Karnataka
2. **Prof. D.K. Srivastava** (*Medical Biochemistry*)
Jamia Hamdard Medical College, New Delhi
3. **Prof. M Sriharibabu** (*General Medicine*) GSL Medical College, Rajahmundry,
Andhra Pradesh
4. **Prof. Pankaj Datta** (*Principal & Prosthodontist*)
Indraprastha Dental College, Ghaziabad

NATIONAL EDITORIAL ADVISORY BOARD

5. **Prof. Samarendra Mahapatro** (*Pediatrician*)
Hi-Tech Medical College, Bhubaneswar, Orissa
6. **Dr. Abhiruchi Galhotra** (*Additional Professor*) Community and Family
Medicine, All India Institute of Medical Sciences, Raipur
7. **Prof. Deepti Pruthvi** (*Pathologist*) SS Institute of Medical Sciences &
Research Center, Davangere, Karnataka
8. **Prof. G S Meena** (*Director Professor*)
Maulana Azad Medical College, New Delhi
9. **Prof. Pradeep Khanna** (*Community Medicine*)
Post Graduate Institute of Medical Sciences, Rohtak, Haryana
10. **Dr. Sunil Mehra** (*Paediatrician & Executive Director*)
MAMTA Health Institute of Mother & Child, New Delhi
11. **Dr Shailendra Handu**, *Associate Professor*, Phrma, DM (Pharma, PGI
Chandigarh)
12. **Dr. A.C. Dhariwal**: *Directorate* of National Vector Borne Disease
Control Programme, Dte. DGHS, Ministry of Health Services, Govt. of
India, Delhi

Print-ISSN: 0976-0245-Electronic-ISSN: 0976-5506, Frequency: Quarterly
(Four issues per volume)

Indian Journal of Public Health Research & Development is a double blind peer reviewed international journal. It deals with all aspects of Public Health including Community Medicine, Public Health, Epidemiology, Occupational Health, Environmental Hazards, Clinical Research, and Public Health Laws and covers all medical specialties concerned with research and development for the masses. The journal strongly encourages reports of research carried out within Indian continent and South East Asia.

The journal has been assigned International Standards Serial Number (ISSN) and is indexed with Index Copernicus (Poland). It is also brought to notice that the journal is being covered by many international databases. The journal is covered by EBSCO (USA), Embase, EMCare & Scopus database. The journal is now part of DST, CSIR, and UGC consortia.

Website : www.ijphrd.com

©All right reserved. The views and opinions expressed are of the authors and not of the Indian Journal of Public Health Research & Development. The journal does not guarantee directly or indirectly the quality or efficacy of any product or service featured in the advertisement in the journal, which are purely commercial.

Editor

Dr. R.K. Sharma

Institute of Medico-legal Publications
Logix Office Tower, Unit No. 1704, Logix City Centre Mall,
Sector- 32, Noida - 201 301 (Uttar Pradesh)

Printed, published and owned by

Dr. R.K. Sharma

Institute of Medico-legal Publications
Logix Office Tower, Unit No. 1704, Logix City Centre Mall,
Sector- 32, Noida - 201 301 (Uttar Pradesh)

Published at

Institute of Medico-legal Publications

Logix Office Tower, Unit No. 1704, Logix City Centre Mall,
Sector- 32, Noida - 201 301 (Uttar Pradesh)



Contents

1. A study to Assess the Knowledge Regarding the Care of Patients on Mechanical Ventilation and Prevention of VAP among Nursing Students of a Selected Nursing College 1
A S Saritha, Jessy Mathew
2. A Study on Job Satisfaction among Employees in Quick Service Restaurants 5
A. Arun, J. Yuvaraj
3. Comparative Evaluation of The Fluoride Release And Rechargability of Chitosan Modified Glass Ionomer Cement and a Glass Ionomer Cement–AN IN VITRO STUDY 9
Binsy Mathew, Abi M Thomas, Rajesh kumar
4. A Study on Determinants of Agricultural Productivity in Tamilnadu 14
C. K. Gomathi, S. N. Sugumar
5. Susceptibility of Gender Entrepreneurship Gap in India–A Revue 18
Chandrachud. S, S. N. Sugumar, S. Thangamayan, S. Sudha
6. Analysis of Medical Tourism and its Economic Impact 21
MS. J. Revathi, S. Jansi Rani
7. Occurrence of Menstrual Irregularities among Adolescent Girls in Selected Area, Dehradun, Uttarakhand . 27
Jyoti Kandpal, Mugdha Devi Sharan Sharma, Upma George
8. Mother’s Knowledge on Nutrition and Incidence of Malnutrition 32
Kalpana Sawane, Sheetal Barde
9. Mortality Pattern amongst Patient Admitted in Tertiary Health Care Center, Rajnandgaon (C.G.) 35
Harshal Mendhe, Kiran Makade, Dhiraj Bhawnani, Daneshwar Singh
10. Effectiveness of Self-Instructional Module on Knowledge Regarding Post-Partum Psychiatric Disorders 41
Leeja Bonny Thomas, Anusha Pradhan
11. A Study to Evaluate the Effectiveness of Structured Teaching Programme on Obesity & its Consequences among Adolescents in Selected Private High Schools in Pune 47
Mangesh V. Jabade, Manu Acha Roy
12. Assessment of the Awareness about Effects of Cell Phone Radiations amongst Students at Symbiosis International (Deemed University) Hill Base Campus 51
Anshika Nikita Singh, Neeti Sharma, Abhay Saraf, Sameer Barve, Yatin Pimple

75. A Study on the Effect of Job Performance on Emotional Labor, Career Turnover Intention, Job Stress, Growth Need 379
Il-Hyun Yun
76. Application of Digital Rubbing Massage in Pain Level, Comfort, and Duration of Labor Phase 385
Sudirman, Sumarni, Hartati, Hendra M., Ismi Rajiani
77. Using Propensity Score Bootstrapping on Determining the Model of the HIV/AIDS Patients' Assistance . 390
Mahdalena, Mahpolah, Ismi Rajiani
78. ARCS Module (*Attention, Relevance, Confidence, Satisfaction*) to Increase Classroom Motivation for Pregnant Women at Public Health Center 395
Agustine Ramie, Mahdalena, Hammad, Ismi Rajiani
79. Interprofessional Education Module in Achieving Ethics/Values, Roles, Responsibilities, Professional Communication Competencies, and Team Collaboration among the College of Health Students 400
Neny Triana, Ismi Rajiani
80. Anxiety Level of Dental Care among Adolescents in Kepulauan Selayar District 403
Lilies Anggarwati Astuti, Nurnaeni, Faiqah Umar, Hasanuddin Tahir, Asmawati Amin
81. Baby Massage With Common Cold Massage Oil on Temperature Change, Pulse Rate, Frequency of Breath, Sleep Quality and Number of Streptococcus Bacteria in Toddlers with Acute Respiratory Infection 407
Melyana Nurul W, Fatatu Malikhah, Kusmini Suprihatin, Sutarmi
82. Risk Factors Affecting Attention Deficit Hyperactivity Disorder Among Early Childhood in the Agricultural Area in Indonesia 411
Istiklaili F, Suwandono A., Suhartono S., Widyorini E., Saputro D.
83. Maternal and Neonatal Outcomes of Elective and Emergency Cesarean Sections 416
Esraa Abdulkareem Mohammed
84. Comorbidities of Phototherapy Used in Neonatal Jaundice in Diyala Governorate, Iraq 422
Saif Hakeem Tofiq, Kareem Assi Obaid, Mazin Razooqi Mohammed
85. Effect of Thyroid Disorder on Liver Function and Some Immunological Parameters 427
Jamela Jouda, Majida G. Maghtoof, Alia Essam Mahmood Alubadi, Youns Atiyah Kamil, Seeham Ali Qasim
86. The Presence of Pathogenic *Leptospira sp* in Water Bodies in Klaten District 433
Novia Tri Astuti, Mateus Sakundarno Adi, Yuliani Setyaningsih, Martini, Lintang Dian Saraswati
87. Dialectic Unity between Threat and Division Sociological Study 438
Majida Shaker Mahdi
88. Job Demands, Low Back Pain, and Job Crafting Behaviors: A Proposed Framework 443
Malek Ahmad Al-Natour, Nor Azimah Chew Abdullah
89. "Educational-Staff Knowledge and Attitude towards Antibiotic Use in Technical Institute of Karbala" . 449
Maytham Salim AL-Nasrawii, Ali abd Al-Latif. G. Mohammed, Mohammad Abdul Baqi Abdul Mohsin, Mohammed A.Merzah

Using Propensity Score Bootstrapping on Determining the Model of the HIV/AIDS Patients' Assistance

Mahdalena¹, Mahpolah¹, Ismi Rajiani²

¹The Ministry of Health Polytechnic Banjarmasin, Indonesia;

²STIA & Port Management Barunawati Surabaya, Indonesia

ABSTRACT

Background: The combination of Propensity Score Stratification (PSS) with a bootstrap method will get an estimate that has better accuracy because the covariates will be more balanced. Propensity score Bootstrapping is used to estimate the response to mentoring HIV/AIDS patients in RSUD dr. H. Moch. Ansari Saleh Banjarmasin South Kalimantan.

Method: The research method used is non-reactive based on secondary data. The study sample was HIV/AIDS patients who had undergone treatment for more than one year at RSUD dr. H. Moch. Ansari Saleh Banjarmasin, South Kalimantan. The response variable is y) CD4 (Cluster of Differentiation) level, while predictor variables (X_j) knowledge, attitude, self-concept, family support, and treatment in the form of antiretroviral therapy.

Results: From the test, the significance of parameters *bootstrap* shows that at the significance level $\alpha = 5\%$ indicating all covariates are significant, namely the covariate of knowledge, attitude, self-concept, family support. Estimation of the PSS bootstrap sample is best in the two strata group with a standard error of 0.0127 and a treatment effect of 0.4175.

Conclusion: The ATE (Average Treatment Effect) value in the univariate response using a proportion test with z test statistics showed that antiretroviral therapy and assistance affected Cluster of Differentiation (CD4) levels.

Keywords: *Propensity Score Stratification, Bootstrap, Antiretroviral Therapy*

INTRODUCTION

Based on reports from the hospital of Dr. H. Moch Ansari Saleh Banjarmasin in 2014, of 422 people who were HIV positive, only 138 people underwent regular therapy, and 14 people dropped out. The number of HIV/AIDS patients who do not undergo treatment regularly, probably because of no companion making HIV/AIDS patients are reluctant to conduct therapy in an orderly manner. As is well known if HIV/AIDS patients do not

take antiretroviral therapy regularly, the HIV cannot be controlled so that it reduces the immune system which will affect CD4 levels, opportunistic infections and the quality of life of HIV/AIDS patients will decrease ⁽¹⁾.

The highest risk of HIV-AIDS transmission in Indonesia comes from sexual relations (62.5%) followed by injecting drugs (16.1%), perinatal (2.7%) and homosexuals (2.4%) Until now; no drugs have been found which can kill the HIV. However, with the discovery of drugs that inhibit the replication of HIV known as Highly Active Antiretroviral Therapy (HAART), the death rate of HIV-AIDS patients can lower ⁽⁵⁾.

The study intends to examine the estimation of propensity score boots on CD4 levels of HIV/AIDS patients who received antiretroviral therapy along with mentoring as a treatment group and who only received antiretroviral treatment as a control group. This causal

Corresponding Author:

Mahdalena

Head of Research and Community Service,
The Ministry of Health Polytechnic Banjarmasin,
Indonesia

Email: lenaf4dl1@gmail.com

research is observational rather than experimental. The estimation method of the propensity score used is the Maximum Likelyhood Estimation (MLE) bootstrapping method. The technique used in the propensity score is a combination stratification method with the method bootstrap to estimate the Average Treatment Effect (ATE) and confounding used only one.

METHOD

This research is a secondary data-based documentation study. To complete the information contained in the research objectives required primary data collection through observational research with a survey approach. Secondary data were obtained from records medical at RSUD dr. H. Moch Ansari Saleh Banjarmasin, covering the characteristics of respondents, antiretroviral therapy, mentoring, CD4 levels. Primary data was also taken through a questionnaire on the respondents include knowledge, attitudes, self-concept, and family support.

The location of this research in Regional General Hospital Dr. H. Moch. Ansari Saleh Banjarmasin Kalimantan Selatan. The research period was 2015 to 2016.

The population of this study was all the status of HIV/AIDS patients in RSUD dr. H. Moch Ansari Saleh Banjarmasin, South Kalimantan, from 2008 to 2016, while the sample is every HIV/AIDS patient, who has undergone treatment for more than six months in dr. H. Moch Ansari Saleh, Banjarmasin, South Kalimantan. After CD4 levels of HIV/AIDS patients were examined for the first time and then examined every six months again.

The variables used in this study consisted of response variables (y)CD4 levels. Predictor variables (x_i) included knowledge, attitudes, self-concept and family support. The treatment variable (Z) as respondents were given antiretroviral therapy and regular and continuous peer assistance. While another variable is the respondents, who were only given antiretroviral therapy without support.

RESULTS

Propensity Score used in this analysis is derived from the following results.

1. Determination of variables confounding using Chi-square: This testing is to find out the relationship between antiretroviral therapy and the

overall predictor variables obtained *value* = 0.000 smaller than $\alpha = 0.05$, meaning there is a relationship between antiretroviral therapy and predictor variables, as well as the relationship between antiretroviral therapy and response variable.

2. Logistic regression model on covariate (x) against confounding (z)

Table 1: Logistic regression test results concurrently with antiretroviral therapy

Antiretroviral Therapy	Chi-square	Significant
Step	86.882	0.000
Block	86.882	0.000
Model	86.882	0.000

Table 1 shows the significance level used is $\alpha = 5$ percent and the test statistics used are G test statistics that follow the distribution Chi-Square. P-value = 0.000 $< \alpha = 0.05$, then the decision taken is that there is at least one predictor variable significant not equal to zero.

Table 2: Logistic regression test results and partial odds ratio antiretroviral therapy

Antiretroviral Therapy	B	Significance	Exp(β)
Knowledge of Good	-2.706	0.000	0.067
Positive Attitudes	-1.754	0.000	0.173
Good Self Concept	-1.112	0.031	0.329
Family Support	-1.232	0.013	0.292
Constant	3.998	0.000	54,511

Table 2 shows that the four variables namely good knowledge, positive attitude, supportive family support have a significant influence on antiretroviral therapy. This result is demonstrated through a p-value less than $\alpha (0.05)$.

Furthermore, the suitability of the model is tested to determine whether there is a difference between the results of the observation and the probability of the prediction. The results of this table are cross-classification of response variables with dichotomous scale. The research shows that the percentage of all observations classified

correctly is 84.0% so that the magnitude of the misclassification is 16.0%. The misclassification value is not too significant, so it can be concluded that the logistic regression model is good enough to classify observations with only antiretroviral therapy (*sensitivity*) percentage of 81.9 percent and the percentage of antiretroviral therapy and assistance (*specificity*) of 85.9 percent.

3. Estimated value of propensity score: The propensity score value in this case, is estimated using a model logistic regression. Therefore the logistic regression model is the same as the *propensity score*, so to know the propensity value, the model parameters must be known in advance. The parameters of the logistic regression model are estimated by the Maximum Likelihood Maximum method. The following is a logistic regression model for antiretroviral therapy :

$$g(x) = 3.998 - 2.706X_{1(1)} - 1.754X_{2(1)} + 1.112X_{3(1)} + 1.232X_{4(1)}$$

Based on the estimation of the parameters above, the obtained *propensity score* will be used at the next stage. The model *propensity score* used is as follows.

$$e(x_i) = \frac{\exp(g(x))}{1 + \exp(g(x))} \dots(1)$$

from equation (1) above, the value of propensity is obtained for 150 research subjects ranging from 0 to 1 because it is a limited opportunity.

4. Estimation of Average Treatment Effect (ATE) or effect of antiretroviral therapy on CD4 levels from HIV/AIDS patients: Providing antiretroviral therapy has a significant effect on CD4 levels in HIV/AIDS patients as p-value is $0.000 < \alpha = 5\%$. The effect of antiretroviral therapy and assistance on increasing CD4 levels in HIV/AIDS patients is 0.678 in the original data.

The bootstrap data giving antiretroviral therapy has a significant effect on CD4 levels of HIV/AIDS patients. It can be seen from the p-value = $0.000 < \alpha = 5\%$. The effect of antiretroviral therapy and accompaniment on increasing CD4 levels in HIV/AIDS patients is 0.690.

Table 3: ATE estimation results for antiretroviral therapy against CD4

Levels of CD4	Sub-Classes	Value of ATE	ATE	Z	p-value
Original	1	0.118421	-0.678876	-11.38	0.000
	2	0.797297			
Bootstrap	1	0.135802	-0.690284	-11.62	0,000
	2	0.826087			

DISCUSSION

Combined antiretroviral (ARV) treatment is the best therapy for human infected patients of Human Immunodeficiency Virus (HIV) to date (2). The primary purpose ARV administration is to reduce the amount of virus (viral load) so that it will improve the HIV patient’s immune status and reduce deaths from opportunistic infections(3). In 2015, according to the World Health Organization (WHO) antiretroviral has been used in 46% of patients HIV in various countries. The use of ARVs has been managed to reduce mortality related to HIV/AIDS from 1.5 million in 2010 to 1.1 million in the year of 2015. Antiretroviral besides being an antiviral is also useful to prevent

transmission of HIV to sexual partners, and HIV transmission from mother to child. Up to finally expected to reduce the number of cases of people new HIV infections in various countries (4).

This study was conducted to determine the effect of antiretroviral therapy on HIV/AIDS patients. The research data contained several covariates involved namely knowledge (X1), attitude (X2), self-concept (X3), family support (X4), and antiretroviral therapy (Z) as well as three response variables (*outcomes*) namely CD4 levels (Y1). Based on the result, it is known that four covariates X (X1, X2, X3, X4) are associated with antiretroviral therapy and are risk factors of CD4 of HIV/AIDS patients.

Logistic regression modeling with more than one predictor variable (simultaneously) needs to be done to examine the multicollinearity relationship between predictor variables. If the connection is very close, it is possible to have predictor variables become insignificant again in testing simultaneously because other predictor variables sufficiently represent it. Concurrent test results showed the significant result.

The principle of propensity score stratification is to divide the subject into several sub-classes based on the propensity score obtained in the previous step. At this stage, the subject is divided into 2 to 5 strata to find strata that prove that the covariates are balanced on all levels and provide an estimate of the standard error value of the smallest Average Treatment Effect (ATE). The distribution of strata was initially carried out on the original data to find the best levels.

Prior estimates with the bootstrap method, first performed at the full data parameter estimation, this is because the criteria for rejection of H_0 using the value *p-value* involving the of the test statistic at complete data. Besides, the significance test of parameters bootstrap shows that at a significance level $\alpha = 5\%$ all covariates are significant, namely the covariate of knowledge, attitude, self-concept and family support.

After their requirements are met, then the next step is to estimate the effect of treatment (ATE) or the impact of antiretroviral therapy on CD4 levels of HIV/AIDS patients. The test was carried out using the Z test, and the results showed that the effect of antiretroviral treatment and accompaniment on increasing CD4 levels in HIV/AIDS patients was 0.678 in the original data.

The *Bootstrap data* giving antiretroviral therapy has a significant effect on CD4 levels of HIV/AIDS patients. The magnitude of the effect is 0.690; thus after bootstrap, it turns out the results are smoother because the fact is stronger.

Based on effect treatment estimates, the provision of antiretroviral therapy and assistance affects increasing CD4 levels. A companion can inspire patients with HIV/AIDS because even though it is incurable but can be extended for life with drugs certain ⁽⁷⁾. Antiretroviral therapy is beneficial in reducing the amount of HIV in the body so that CD4 (lymphocytes *cluster of differentiation 4*) can be prevented. After the administration of antiretroviral therapy drugs for six months can usually

be achieved the number of undetectable viruses and the number of CD4 lymphocytes increases ⁽⁸⁾. As a result of the risk of opportunistic infections decreases and the quality of life of patients increases ⁽⁹⁾.

HIV/AIDS patients who are given antiretroviral therapy and mentoring will have an increase in CD4 levels. Through the need to increase knowledge, a positive attitude is formed which makes the concept of self-good. No less important for the family to support their family support HIV/AIDS patients with HIV feel cared for and being considered in the family.

The lower CD4 count identifies a decreased immune system so that pathogens that cause infection can enter the body together. Giving ARVs can increase CD4 levels, which also means rising immunity to prevent opportunistic infections⁽⁶⁾.

CONCLUSION

From the research, it can be obtained estimator propensity score stratification bootstrap. Estimated PSS bootstrap samples are best in the two levels group. The effect of antiretroviral therapy and accompaniment to CD4 levels significantly impact on the HIV patient.

Ethical Clearance: Ethical clearance was obtained from the the Health Research Ethics Committee at the Faculty of Public Health Airlangga University with registration number 161-KEPK dated 14 April 2016. We also wish to thank all the participants who contributed to this study.

Conflict of Interest: Nil.

Source of Funding: The Ministry of Health Polytechnic Banjarmasin, Indonesia.

REFERENCES

1. Aggarwal S, Yu L, Hasjim B, Lee DH, Kim E, Lee JB, Lee K, Tse C, Anshus A, Yu AR, Louthan M. Stigma and negative self-perceptions of young people living with human immunodeficiency virus in Bandung, Indonesia: a case series. *International health*. 2018 Jun 12.
2. Austin PC, Mamdani MM. A comparison of propensity score methods: a case-study is estimating the effectiveness of post-AMI statin use. *Statistics in medicine*. 2006 Jun 30;25(12):2084-106.

3. Amon JJ, Todrys KW. Fear of foreigners: HIV-related restrictions on entry, stay, and residence. *Journal of the International AIDS Society*. 2008 Dec;11(1):8.
4. Liping M, Peng X, Haijiang L, Lahong J, Fan L. Quality of life of people living with HIV/AIDS: a cross-sectional study in Zhejiang province, China. *PloS one*. 2015 Aug 26;10(8):e0135705.
5. Pohan HT. Opportunistic infection of HIV-infected/AIDS patients in Indonesia: problems and challenge. *Acta Med Indonesia*. 2006;38(1):135-51.
6. Bluthenthal RN, Palar K, Mendel P, Kanouse DE, Corbin DE, Derose KP. Attitudes and beliefs related to HIV/AIDS in urban religious congregations: Barriers and opportunities for HIV-related interventions. *Social science & medicine*. 2012 May 1;74(10):1520-7.
7. Butt L. Local biologies and HIV/AIDS in highlands Papua, Indonesia. *Culture, Medicine, and Psychiatry*. 2013 Mar 1;37(1):179-94.
8. Ho DD, Neumann AU, Perelson AS, Chen W, Leonard JM, Markowitz M. Rapid turnover of plasma virions and CD4 lymphocytes in HIV-1 infection. *Nature*. 1995 Jan;373(6510):123.
9. Dalgleish AG, Beverley PC, Clapham PR, Crawford DH, Greaves MF, Weiss RA. The CD4 (T4) antigen is an essential component of the receptor for the AIDS retrovirus. *Nature*. 1984 Dec;312(5996):763.